

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 5/04, 5/06, 1/16, 1/20, 15/67, 15/81, 15/82, 15/90, A01H 1/00, 5/00		A1	(11) International Publication Number: WO 00/52144
			(43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/US00/05315		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 February 2000 (28.02.00)			
(30) Priority Data: 09/261,825 3 March 1999 (03.03.99) US			
(71) Applicant: UNIVERSITY OF TEXAS [US/US]; 201 West 7th Street, Austin, TX 78701-2981 (US).			
(72) Inventors: THOMAS, Collin, E.; 8702, Parkfield Drive, Austin, TX 78758 (US). WINDSOR, J., Brian; 4701 Staggerbrush Road, Apt. 636, Austin, TX 78749 (US). ROUX, Stan, J.; 8510 Forest Heights Lane, Austin, TX 78749-3505 (US). LLOYD, Alan, M.; 2505 Forest Bend Drive, Austin, TX 78704 (US). HURLEY, Laurence; 5811 Mesa Drive, Apt. 1316, Austin, TX 78731 (US).			
(74) Agents: MCDANIEL, C., Steven et al.; McDaniel & Associates, P.C., P.O. Box 2244, Austin, TX 78768-2244 (US).		Published <i>With international search report.</i>	
(54) Title: GENETIC AND EPIGENETIC MANIPULATION OF ABC TRANSPORTERS AND ECTO-PHOSPHATASES FOR THE CONFERENCE OF DRUG RESISTANCE AND FOR THE LOSS OF DRUG RESISTANCE IN BIOLOGICAL SYSTEMS AND METHODS FOR THE DETECTION OF ECTO-PHOSPHATASE INHIBITORS			
(57) Abstract <p>The present invention relates to methods for modulating the resistance of cells to foreign compounds, i.e. drugs, antibiotics, etc by altering the ATP gradient across biological membranes. The altering of the ATP gradient across biological membranes is achieved through the manipulation of ecto-phosphatase activity and ABC transporter molecule activity which may be useful to confer herbicide resistance to plants, confer antibiotic resistance to bacteria, confer drug resistance to yeast cells, or to reduce resistance in cells to facilitate chemotherapeutic treatments, and to reduce resistance in bacteria and yeast. The present invention is also directed to the methods for identifying ecto-phosphatase inhibitors and uses thereof.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

GENETIC AND EPIGENETIC MANIPULATION OF ABC TRANSPORTERS AND
ECTO-PHOSPHATASES FOR THE CONFERENCE OF DRUG RESISTANCE AND FOR
THE LOSS OF DRUG RESISTANCE IN BIOLOGICAL SYSTEMS AND METHODS
FOR THE DETECTION OF ECTO-PHOSPHATASE INHIBITORS

1. INTRODUCTION

The present invention is concerned with modulating the drug resistance pathways of cells in order to either confer or overcome resistance to certain drug molecules. Such modulation entails modulation of an extra-cellular phosphatase (ecto-phosphatase) and an ABC (ATP-binding cassette) transporter in order to achieve the desired effect on drug resistance. Stimulation of the ecto-phosphatase either alone or together with stimulation of the ABC transporter yields an increased resistance to drug molecules while inhibition of the ecto-phosphatase alone or together with the ABC transporter yields reduced resistance to the drug molecule. Drug resistance is achieved through the altering of the ATP gradient across biological membranes which is effectuated through the modulation of an ecto-phosphatase either alone or together with an ABC transporter molecule. Modulation of drug resistance as described herein is useful in conferring herbicide resistance to plants; conferring drug resistance to microorganisms and tissue culture cells; reducing drug resistance in tumor cells for improved chemotherapy applications; and reducing resistance to antibiotics, antifungal agents, and other drugs in microorganisms for the treatment of infections and disease. The present invention is also directed to methods for identifying inhibitors of ecto-phosphatases and uses thereof.

BACKGROUND OF THE INVENTION

Transport Processes

Cells can use a phenomenon called symport to move soluble products across biological membranes. Symport is a form of coupled movement of two solutes in the same direction across a membrane by a single carrier. Examples of proton and sodium-linked symport systems are found in nearly all living systems. The energetics of the transport event depend on the relative size and electrical nature of the gradient of solutes.

Transport processes have been classified on the basis of their energy-coupling mechanisms. Currently there are four classifications: (1) Primary Active Transport which uses either a chemical, light or electrical energy source, (2) Group Translocation which uses

chemical energy sources, (3) Secondary Active Transport which uses either a sodium or proton electrochemical gradient energy source, and (4) Facilitated Diffusion which does not require an energy source. Meyers, R. A., 1997, *Encyclopedia of Molecular Biology and Molecular Medicine* 6:125-133. The present invention is related to transport molecules
5 belonging to the first class of transport processes, primary active transport, and therefore, this type of transport will be discussed in further detail.

Primary active transport refers to a process whereby a "primary" source of energy is used to drive the active accumulation of a solute into or extrusion of a solute from a cell. Transport proteins include P-type ATPases and ABC-type ATPases. These types of
10 transport systems are found in both eukaryotes and prokaryotes. The bacterial ABC-type transporters, which are ATP-driven solute pumps, have eukaryotic counterparts. Additionally, many transmembrane solute transport proteins exhibit a common structural motif. The proteins in these families consist of units or domains that pass through the membrane six times, each time as an α -helix. This has led to the suggestion that many transport proteins
15 share a common evolutionary origin, but this is not true of several distinct families of transport proteins. Numerous structurally distinct bacterial permeases, as well as several homologous eukaryotic transport systems, share a common organization. Meyers, R. A., 1997, *Encyclopedia of Molecular Biology and Molecular Medicine* 6:125-133. Two hydrophilic domains or proteins function to couple ATP hydrolysis in the cytoplasm to activate substrate
20 uptake or efflux, and two hydrophobic domains or proteins function as the transmembrane substrate channels. These proteins or protein domains constitute what is referred to as the ABC (ATP-binding cassette) superfamily. Either the two hydrophilic domains or proteins or the two hydrophobic domains or proteins (or both) may exist either as heterodimers or homodimers. If, as in most bacterial systems, each of these constituents is a distinct protein,
25 then either two, three, or four genes will code for them, depending on whether both are homodimers, one is a homodimer and one is a heterodimer, or both are heterodimers, respectively. The best characterized of the eukaryotic proteins included in this family are the multidrug-resistance (MDR) transporter and the cystic fibrosis related chloride ion channel of mammalian cells (cystic fibrosis transmembrane conductance regulator or CFTR). Meyers, R.
30 A., 1997, *Encyclopedia of Molecular Biology and Molecular Medicine* 6:125-133.

Multidrug Resistance

Multidrug resistance (MDR) is a general term that refers to the phenotype of cells or microorganisms that exhibit resistance to different, chemically dissimilar, cytotoxic compounds. MDR can develop after sequential or simultaneous exposure to various drugs. MDR can also develop before exposure to many compounds to which a cell or microorganism may be found to be resistant. MDR which develops before exposure is frequently due to a genetic event which causes the altered expression and/or mutation of an ATP-binding cassette (ABC) transporter. Wadkins, R. M. and Roepe, P. D., 1997, *International Review of Cytology* 171:121-165. This is true for both eukaryotes and prokaryotes. *Id.*

One prominent member of the ABC family, P-glycoprotein (Pgp; also known as multidrug resistance protein or MDR1), which is a plasma-membrane glycoprotein that confers a multidrug resistance (MDR) phenotype on cells, is of considerable interest because it provides one mechanism of possibly inhibiting resistance in tumor cells to chemotherapeutic agents. Senior, A E. et al., 1995, *FEBS Letters* 377:285-289. Pgp is a single polypeptide of ~1280 amino acids with the typical ABC transporter structure profile. Studies have shown that over-expression of Pgp is responsible for the ATP-dependent extrusion of a variety of compounds, including chemotherapeutic drugs, from cells. Abraham, E. H. et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:312-316.

Over one-hundred ABC transporters have been identified in species ranging from *Escherichia coli* to humans. Higgins C. F., 1995, *Cell* 82:693-696. For example, the bacteria *Lactococcus lactis* expresses an ABC transporter, LmrA, which mediates antibiotic resistance by extruding amphiphilic compounds from the inner leaflet of the cytoplasmic membrane. van Veen H.W. et al., 1998, *Nature* 391:291-295. Furthermore, over-expression of LmrA can confer MDR in human lung fibroblasts and LmrA has similar molecular and biochemical properties to Pgp. *Id.* This demonstrates that bacterial LmrA and Pgp are functionally interchangeable. *Id.* Additionally, the plant *Arabidopsis thaliana* encodes an ATP transporter, AtPGP-1, which is a putative Pgp homolog. Dudler, R. and Hertig, C., 1992, *Journal of Biological Chemistry* 267:5882-5888. Similarly, the yeast *Saccharomyces cerevisiae* equivalent of Pgp, STS1 (Bissinger, P.H and Kucher, K., 1994, *J. Biol. Chem.* 269:4180-4186), has been cloned and shown to confer multidrug resistance when over-

expressed in yeast, as has the yeast Pdr5p (Kolaczkowski et al., 1996, *J. Biol. Chem.* 271:31543-31548). Taken together, these results suggest that this type of multidrug resistance efflux pump is conserved from bacteria to humans.

While various theories of ABC transporter function have become popular, there is still no precise molecular-level description for the mechanism by which over-expression lowers intracellular accumulation of drugs, in particular how Pgp lowers intracellular accumulation of chemotherapeutic drugs. However, it has been shown that Pgp over-expression also changes plasma membrane electrical potential and intracellular pH which could potentially greatly affect the cellular flux of a large number of compounds to which Pgp confers resistance. Randy M. Wadkins and Paul D. Roepe, 1997, *International Review of Cytology* 171:121-165. Also included in the ABC transporter superfamily are the Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) and the Sulfonyl Urea Receptor (SUR). CFTR and SUR are expressed in the lung epithelium and the β cells of the pancreas, respectively, as well as in other tissues. CFTR functions as a low conductance ATP and cyclic AMP-dependent Cl^- channel that also appears to have additional important functions, such as modulation of epithelial Na^+ conductance and regulation of outwardly rectified chloride channels. Wadkins, R. M. and Roepe, P.D., 1997, *International Review of Cytology* 171:121-165. Mutations in the CFTR gene produce altered CFTR proteins with defects in CFTR function, leading to profound alterations in epithelial salt transport and altered mucous properties in cystic fibrosis patients that result in chronic lung infections associated with the disease. *Id.* SUR is triggered by sulfonyl urea drugs to depolarize pancreatic β cells that leads to Ca^{2+} influx, which stimulates fusion of insulin-containing vesicles to the plasma membrane. *Id.* An ATP transporter hypothesis has been suggested for Pgp, CFTR and SUR which theorizes that these ABC transporters function as ATP transport channels. Abraham, E. H. et al, 1993, *Proc. Natl. Acad. Sci. USA* 90:312-316; Schweibert, E. M., 1995, *Cell* 81:1063-1073; and Al-Awqati, Q., 1995, *Science* 269:805-806. The ATP channel hypothesis, however, has been viewed with skepticism. This is partly due to the inability to show the same results with preparations including purified and reconstituted CFTR, suggesting that the ATP conductance that was originally observed may have been mediated by another protein, not present in the purified system, that is influenced by CFTR. Wadkins, R.

M. and Roepe, P. D., 1997, *International Review of Cytology* 171:121-165. There has been no such negative data reported with respect to the ATP channel hypothesis for Pgp or SUR, but the controversy over CFTR has raised doubt for Pgp and SUR as well.

In support of the ATP channel hypothesis, Huang et al. (*Biochem. Biophys. Res. Commun.* 182:836-843 (1992)) have suggested that extracellular ATP leads to elevations in pH, and Weiner et al. (*J. Biol. Chem.* 261:4529-4534 (1986)) have suggested that extracellular ATP may regulate Na^+/H^+ exchange in Ehrlich ascites tumor cells. It has also been observed that changes in Pgp levels affects pH and plasma membrane electrical potentials which could be connected to recent observations suggesting the involvement of ATP transport in MDR.

Additionally, Abraham et al. (*Proc. Natl. Acad. Sci. USA* 90:312-316 (1993)) have reported that the addition of extracellular ATP to MDR cell lines confers sensitivity to drugs abolishing MDR. The data for this effect were not presented in the article and no further explanation was given for this phenomenon. Furthermore, there have been no subsequent publications addressing or explaining this effect.

Furthermore, Ujhazy et al. (*Int. J. Cancer* 68:493-500 (1996)) have shown that ecto-5'-nucleotidase is up-regulated in certain MDR cell lines. Ecto-5'-nucleotidase is the final enzyme in the extracellular pathway for salvage of adenosine from phosphorylated purines. Zimmerman H., 1992, *Biochem. J.* 285:345-365. The proposed hypothesis for the involvement of ecto-5'-nucleotidase in drug resistance considers its role in the maintenance of intracellular ATP pools through the adenosine salvage pathway. Ujhazy et al., 1996, *Int. J. Cancer* 68:493-500. Ecto-5'-nucleotidase specifically acts in adenosine salvage pathways, converting AMP to adenosine which is more readily taken up by the cell and utilized as a precursor for ATP production. Therefore, ecto-5'-nucleotidase may be acting in certain MDR cell lines as a mechanism by which the cell circumvents the loss of ATP (due to up-regulated transport proteins which possibly form ATP transport channels) by creating higher levels of adenosine from which the cell can produce ATP. Correspondingly, 63% of MDR cell line variants tested expressed ecto-5'- nucleotidase. These observations suggested that a salvage mechanism for extracellular nucleotides may be another way by which certain MDR cells counterbalance their ATP losses from efflux induced by the over-expression of ABC transporters involved in MDR. Consistent with this hypothesis, inhibitors of ecto-5'-

nucleotidase conferred sensitivity to certain drugs in MDR cell lines which over-express the ecto-5'-nucleotidase.

It is also interesting to note that yeast, which do not have an adenosine salvage pathway (Boyum, R. and Guidotti, G., 1997, *Microbiology* 143:1901-1908), do contain a Pgp-like gene called STS1 (Bissinger, P.H. and Kucher, K., 1994, *J. Biol. Chem.* 269:4180-4186. Therefore, since the adenosine salvage pathway is unlikely to be involved in yeast multidrug resistance, other mechanisms are likely to exist.

Recent reports have confirmed the existence of ATP in the extracellular matrix (ECM) of both multicellular organisms and unicellular organisms. Sedaa, K. et al., 1990, *J. Pharmacol. Exp. Ther.* 252:1060-1067 and Boyum, R. and Guidotti, G., 1997, *Microbiology* 143:1901-1908, respectively. However, no such reports are available which suggest the existence of ATP in the ECM of plants before the present invention. These reports have prompted further investigations of the fate of ATP outside the cell. One of the largest gradients in biological systems is that of ATP. It is a million-fold more concentrated inside the cell than outside. Apyrases are enzymes whose unifying characteristic is their ability to hydrolyze the gamma phosphate of ATP and to a lesser extent, the beta phosphate of ADP. Plesner, L., 1995, *Int. Rev. Cytol.* 158:141-214. Most apyrases are expressed as plasma membrane associated proteins with their hydrolytic activity facing into the ECM. Wang, T. and Guidotti, G., 1996, *J. Biol. Chem.* 271:9898-9901. Extracellular apyrases are generally referred to as ecto-apyrases. Given reports that show the existence of extracellular ATP, one observation regarding ecto-apyrase is that it hydrolyzes the extracellular ATP. In fact, work in animal systems has shown that apyrases hydrolyze ATP in the ECM as part of the adenosine salvage pathway con-jointly with ecto-5' ectonucleotidase. Che, M., 1992, *J. Biol. Chem.* 267:9684-9688. The existence of a similar ecto-apyrase system has not been reported in plants prior to the present invention. Additionally, ecto-apyrases have not been shown, prior to the present invention, to have a role in MDR.

While some references appear to indicate that MDR may act at the level of ATP transport, the role of ATP in MDR has not been adequately elucidated and has remained a point of contention in the field. The present invention provides insight into the role of ATP transport in MDR by showing that the extracellular ATP pool in cells is critical in MDR. While the adenosine salvage pathway may help compensate for ATP losses in MDR by providing a mechanism to recoup adenosine, it is not the critical aspect of the role of ATP in

MDR as evidenced by the observation that only a subset of MDR cell lines resort to this mechanism via the up-regulation of ecto-5'-nucleotidase to maintain drug resistance. In fact, the previous data teach away from modulating extracellular ATP levels and place the focus on mechanisms which are involved in modulating intracellular ATP levels. Since AMP is the preferred substrate for ecto-5'-nucleotidase, with ATP and ADP being poor substrates (Zimmerman, H., 1992, *Biochem. J.* 285:345-365), it is unlikely that ecto-5'-nucleotidase is involved in modulating extracellular levels of ATP. While high levels of ATP have been demonstrated to be useful in the inhibition of tumor growth, its effects on tumor cells have been shown to prevent cell growth and induce cell death through the inhibition of the S phase of the cell cycle. U.S. Patent 4,880,918. There has been no implication, prior to the present invention, of the importance of modulating extracellular ATP levels in MDR.

Additionally, there has been no identification of specific inhibitors of apyrase (an ecto-phosphatase). Such inhibitors and methods for identifying such inhibitors would be useful for studying the importance of ecto-phosphatases in MDR, for modulating MDR and in industrial applications (*e.g.* determining the titer of microbia in soil).

It would be particularly useful to have more effective mechanisms by which to modulate drug resistance in various organisms. In particular, since the use of Pgp inhibitors has not been totally efficient in overcoming the resistance seen in tumor cells which have been repeatedly exposed to chemotherapeutic agents, it would be useful to have other mechanisms by which to combat such resistance in tumor cells to provide more effective chemotherapeutic treatments. Furthermore, there are many other applications for the modulation of drug resistance which are contemplated by the present invention, such as the engineering of herbicide resistant plants for use in agriculture.

SUMMARY OF THE INVENTION

The present invention is directed to a method for the modulation of drug resistance in cells. In one embodiment, resistance is conferred through over-expression by genetic manipulation of ABC transporters and ecto-phosphatases which are capable of affecting extracellular ATP pools and thus affecting the ATP gradient across biological membranes. Conferred resistance is useful to achieve herbicide resistance in plants, drug resistance in yeast (*i.e.* resistance to anti-fungal agents) in biotechnology applications, antibiotic resistance in bacteria in biotechnology applications and for drug resistance in

eukaryotic tissue culture cells in biotechnology applications. In another embodiment, loss of drug resistance is achieved by suppressing the breakdown of extracellular ATP through the down-regulation of ecto-phosphatases in the presence or absence of the down-regulation of ABC transporters. Loss of resistance is useful to mitigate drug resistance problems associated with chemotherapy and in the treatment of infections from resistant strains of microorganisms. The modulation of drug resistance is achieved, at least in part, by altering the ATP gradient across biological membranes through the aforementioned manipulation of ABC transporters and ecto-phosphatases. The present invention is also directed to methods for the identification of inhibitors of ecto-phosphatases and uses thereof.

BRIEF DESCRIPTION OF THE FIGURES.

FIGURE 1. Expression of apyrase in pea and in transgenic plants (A) Immunoblot analysis of subcellular fractions from etiolated pea plants. (B) Top, the total phosphate accumulated in the shoots of three independent transgenic plants. Bottom, a corresponding immunoblot performed on protein from ECM of wild-type and transgenic plants. (C) Assay of phosphatase activity in the ECM fraction of OE1 and wild-type.

FIGURE 2. Transport of the products of ATP hydrolysis by transgenic plants overexpressing apyrase and by wild-type plants.

FIGURE 3. Conference of resistance to cycloheximide (A and B) and nigericin (C and D) in wild-type and ecto-phosphatase deficient yeast over-expressing the *Arabidopsis* plant ABC transporter, AtPGP-1.

FIGURE 4. Conference of resistance to cycloheximide (A) and cytokinin (B) in *Arabidopsis* plants over-expressing either the ecto-phosphatase, apyrase, or the ABC transporter, AtPGP-1.

FIGURE 5. Graph showing the growth turbidity of YMR4 yeast over-expressing the *Arabidopsis* plant ABC transporter AtPGP-1 grown in cycloheximide (A) or nigericin (B).

FIGURE 6. Graph showing germination rate of *Arabidopsis* plants grown in the presence of cycloheximide which over-express either the ecto-phosphatase, apyrase, or the ABC transporter AtPGP-1.

FIGURE 7. Graph of steady-state levels of ATP in the extracellular fluid of wild-type yeast cells grown in the presence or absence of glucose and in the presence or absence of over-expression of the *Arabidopsis* plant ABC transporter, AtPGP-1.

5 FIGURE 8. Graph showing that over-expression of *Arabidopsis* plant ABC transporter, AtPGP-1, in yeast can double the steady-state levels of ATP in the extracellular fluid.

FIGURE 9. Graph showing that a yeast mutant, YMR4, that has a deficient ecto-phosphatase, accumulates ATP in the extracellular fluid and the over-expression of AtPGP-1 increases the accumulation of ATP.

10 FIGURE 10. Graph showing results of a pulse-chase experiment in either wild-type yeast cells or a yeast mutant, YMR4, which is deficient in ecto-phosphatase activity, in the presence and absence of over-expression of *Arabidopsis* plant ABC transporter, AtPGP-1, demonstrating an early differential ATP efflux of cells over-expressing AtPGP-1.

15 FIGURE 11. Graph of ATP levels on the surface of leaves of *Arabidopsis* plants over-expressing AtPGP-1 (MDR1).

FIGURE 12. Effects of phosphatase inhibitor in wild-type and AtPGP-1 (MDR1) overexpressing *Arabidopsis* plants.

DETAILED DESCRIPTION OF THE INVENTION

20 For purposes of clarity of description, and not by way of limitation, the detailed description of the invention is divided into the following subsections:

- (i) conference of herbicide resistance in plants;
- (ii) conference of drug resistance in recombinant research applications;
- (iii) inhibition of drug resistance in chemotherapy;
- 25 (iv) inhibition of drug resistance in microorganisms to treat infection;
- (v) ecto-phosphatase inhibition

Conference of herbicide resistance in plants

30 The present invention is directed to a method for the modulation of drug resistance in plants, particularly herbicide resistance, in part through the manipulation of the ATP gradient across biological membranes. In accordance with the invention, the manipulation of extracellular ATP levels and hence the ATP gradient across biological membranes in plant cells by the over-expression of a MDR-ABC transporter and an ecto-

phosphatase, results in resistance to certain plant hormones, drugs and herbicides. Such resistance is useful in horticulture of recombinant crops for the elimination of other unwanted plants (e.g. weeds) which are not resistant. The invention is based, in part, on the unexpected observation that the over-expression of either an ecto-phosphatase, or an ABC transporter can confer resistance to certain drugs and herbicides in plants.

Up-regulation as used herein refers to increasing the activity of a molecule within a cell by either providing an outside source of the molecule (e.g. an expression cassette containing a DNA encoding the molecule) either in single copy or multiple copies which when expressed in the cell increases the amount of the molecule in the cell, by increasing the transcription of the endogenous or exogenous molecule to increase the amount of the molecule in the cell, or by modifying the exogenous or endogenous molecule in the cell post-translationally to achieve an increase in activity of the molecule. Down-regulation as used herein refers to decreasing the activity of a molecule in a cell by either decreasing the amount of the molecule in the cell (this may be achieved by over-expression of an anti-sense RNA corresponding to the molecule or by inhibiting factors necessary for the expression of the molecule) or by modifying the exogenous or endogenous molecule in the cell post-translationally to achieve a decrease in activity. Such post translational modifications may include phosphorylation, adenylation, glycosylation, ubiquitinylation, acetylation, methylation, farnesylation, myristilation and sulfation.

The ecto-phosphatases remove phosphate from any ATP extruded from the cell, rendering the ATP ineffectual for transport of drugs back into the cell. Ecto-phosphatases as referred to herein do not include extracellular phosphatases involved in the adenosine salvage pathway. MDR ABC transporters form channels which facilitate the efflux of molecules, including drugs, from cells. This efflux is likely effectuated through the "piggy-back" efflux of drug molecules with ATP, a phenomenon known as symport.

In one embodiment of the invention, the over-expression of an ecto-phosphatase confers drug resistance in both wild-type and/or genetically engineered plants. This effect is seen in plant cells over-expressing plant apyrase grown in the presence of (1) cycloheximide, a potent inhibitor of protein expression, (2) nigericin, an antibiotic which effects ion transport, and (3) N₆ (2-isopentenyl) adenine, a cytokinin plant hormone which is herbicidal at micromolar and millimolar concentrations.

In another embodiment of the invention, the over-expression of an ABC transporter confers drug resistance in wild-type and genetically engineered plants. In a preferred embodiment, the ABC transporter which is over-expressed is the *Arabidopsis* ABC transporter AtPGP-1. The over-expression of AtPGP-1 can confer resistance in plants to cycloheximide, nigericin and cytokinins.

In a preferred embodiment of the invention the effect of over-expression of both an MDR-ABC transporter and an ecto-phosphatase is enhancement of the ATP gradient across biological membranes and thus stimulation of resistance to certain plant hormones and herbicides. In a particularly preferred embodiment of the invention, the MDR-ABC transporter which is over-expressed is the *Arabidopsis* AtPGP-1 and the ecto-phosphatase that is over-expressed is apyrase.

The invention particularly contemplates the conference of resistance in plants to herbicides which resemble established drugs implicated in multidrug resistance, as well as plant hormones such as cytokinin, auxins, gibberellins and brassinosteroids. The present invention also contemplates the conference of resistance in plants to the nonlimiting list of chemicals, such as those set forth in Table 1*.

TABLE 1

Common Name	Chemical Name
acetochlor	-chloro-N-(ethoxymethyl)-N-(2-ethyl-6-methylphenyl)acetamide
acifluorfen	5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoic acid
acrolein	2-propenal
alachlor	2-chloro-N-(2,6-diethylphenyl)-N-(methoxymethyl)acetamide
allyl alcohol	2-propen-1-ol
ametryn	N-ethyl-N'-(1-methylethyl)-6-(methylthio)-1,3,5-triazine-2,4-diamine
amitrole	1H-1,2,4-triazol-3-amine
AMS	ammonium sulfamate
arsenic acid	arsenic acid
asulam	methyl[(4-aminophenyl)sulfonyl]carbamate
atraton	N-ethyl-6-methoxy-N'-(1-methylethyl)-1,3,5-triazine-2,4-diamine
atrazine	6-chloro-N-ethyl-N'-(1-methylethyl)-1,3,5-triazine-2,4-diamine

	azafenidin	2-[2,4-dichloro-5-(2-propynyloxy)phenyl]-5,6,7,8-tetrahydro-1,2,4-triazolo[4,3-a]pyridin-3(2H)-one
	azimsulfuron	N-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-1-methyl-4-(2-methyl-2H-tetrazol-5-yl)-1H-pyrazole-5-sulfonamide
	barban	4-chloro-2-butynyl 3-chlorophenylcarbamate
	BCPC	1-methylpropyl 3-chlorophenylcarbamate
5	benazolin	4-chloro-2-oxo-3(2H)-benzothiazoleacetic acid
	benefin	N-butyl-N-ethyl-2,6-dinitro-4-(trifluoromethyl)benzenamine
	bensulfuron	2-[[[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]methyl]benzoic acid
	bensulide	O,O-bis(1-methylethyl)S-[2-[(phenylsulfonyl)amino]ethyl]phosphorodithioate
	bentazon	3-(1-methylethyl)-(1H)-2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide
10	benzadox	[(benzoylamino)oxy]acetic acid
	benzipram	3,5-dimethyl-N-(1-methylethyl)-N-(phenylmethyl)benzamide
	benzofluor	N-[4-(ethylthio)-2-(trifluoromethyl)phenyl]methanesulfonamide
	benzoylprop	N-benzoyl-N-(3,4-dichlorophenyl)-DL-alanine
	benzthiazuron	N-2-benzothiazolyl-N'-methylurea
15	bifenox	methyl 5-(2,4-dichlorophenoxy)-2-nitrobenzoate
	borax	sodium tetraborate
	bromacil	5-bromo-6-methyl-3-(1-methylpropyl)-2,4(1H, 3H)pyrimidinedione
	bromofenoxim	3,5-dibromo-4-hydroxybenzaldehyde O-(2,4-dinitrophenyl)oxime
	bromoxynil	3,5-dibromo-4-hydroxybenzonitrile
20	butachlor	N-(butoxymethyl)-2-chloro-N-(2,6-diethylphenyl)acetamide
	butam	2,2-dimethyl-N-(1-methylethyl)-N-(phenylmethyl)propanamide
	butamifos	O-ethyl O-(5-methyl-2-nitrophenyl)1-methylpropylphosphoramidothioate
	buthidazole	3-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-4-hydroxy-1-methyl-2-imidazolidinone
	butralin	4-(1,1-dimethylethyl)-N-(1-methylpropyl)-2,6-dinitrobenzenamine
25	buturon	N'-(4-chlorophenyl)-N-methyl-N-(1-methyl-2-propynyl)urea
	butylate	S-ethyl bis(2-methylpropyl)carbamothioate

	cacodylic acid	dimethyl arsinic acid
	cambendichlor	(phenylimino)di-2,1-ethanediyl bis(3,6-dichloro-2-methoxybenzoate)
	carbetamide	N-ethyl-2-[[[(phenylamino)carbonyl]oxy]propanamide (R)-isomer
	CDAA	2-chloro-N,N-di-2-propenylacetamide
5	carfentrazone	"the alpha character" 2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoic acid
	CDEA	2-chloro-N,N-diethylacetamide
	CDEC	2-chloro-2-propenyl diethylcarbamodithioate
	CEPC	2-chloroethyl (3-chlorophenyl)carbamate
	chloramben	3-amino-2,5-dichlorobenzoic acid
10	chlorazine	6-chloro-N,N,N',N'-tetraethyl-1,3,5-triazine-2,4-diamine
	chlorbromuron	N'-(4-bromo-3-chlorophenyl)-N-methoxy-N-methylurea
	chlorbufam	1-methyl-2-propynyl (3-chlorophenyl)carbamate
	chlorfluorenol	2-chloro-9-hydroxy-9H-fluorene-9-carboxylic acid
	chlorimuron	2-[[[(4-chloro-6-methoxy-2-pyrimidinyl)amino]carbonyl]amino[sulfonyl]benzoic acid
15	chloroxuron	N'-(4-(4-chlorophenoxy)phenyl)-N,N-dimethylurea
	chlorpropham	1-methylethyl 3-chlorophenylcarbamate
	chlorsulfuron	2-chloro-N-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide
	chlorthiamid	2,6-dichlorobenzenecarbothiamide
	chlortoluron	N'-(3-chloro-4-methylphenyl)-N,N-dimethylurea
20	cinmethylin	exo-(±)-1-methyl-4-(1-methylethyl)-2-[(2-methylphenyl)methoxy]-7-oxabicyclo[2.2.1]heptane
	cisanilide	cis-2,5-dimethyl-N-phenyl-1-pyrrolidinecarboxamide
	clethodim	(E,E)-(±)-2-[1-[[[(3-chloro-2-propenyl)oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one
	clofop	2-[4-(4-chlorophenoxy)phenoxy]propanoic acid
	clomazone	2-[(2-chlorophenyl)methyl]-4,4-dimethyl-3-isoxazolidinone
25	cloproxydim	(E,E)-2-[1-[[[(3-chloro-2-propenyl)oxy]imino]butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one

	cloransulam	3-chloro-2-[[[(5-ethoxy-7-fluoro[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)sulfonyl]amino]benzoic acid
	clopyralid	3,6-dichloro-2-pyridinecarboxylic acid
	CMA	calcium salt of MAA
	copper sulfate	copper sulfate
5	4-CPA	(4-chlorophenoxy)acetic acid
	4-CPB	4-(4-chlorophenoxy)butyric acid
	CPMF	1-chloro-N'-(3,4-dichlorophenyl)-N,N-dimethylformamidine
	4-CPP	2-(4-chlorophenoxy)propionic acid
	CPPC	2-chloro-1-methylethyl (3-chlorophenyl)carbamate
10	cyanazine	2-[[[4-chloro-6-(ethylamino)-1,3,5-triazin-2-yl]amino]-2-methylpropanenitrile
	cycloate	S-ethyl cyclohexylethylcarbamothioate
	cyclosulfamuron	N-[[[2-(cyclopropylcarbonyl)phenyl]amino]sulfonyl]-N'-(4,6-dimethoxy-2-pyrimidinyl)urea
	cycluron	N'-cyclooctyl-N,N-dimethylurea
	cyhalofop	(R)-2-[4-(4-cyano-2-fluorophenoxy)phenoxy]propanoic acid
15	cyperquat	1-methyl-4-phenylpyridinium
	cyprazine	6-chloro-N-cyclopropyl-N'-(1-methylethyl)-1,3,5-triazine-2,4-diamine
	cyprazole	N-[5-(2-chloro-1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]cyclopropanecarboxamide
	cypromid	N-(3,4-dichlorophenyl)cyclopropanecarboxamide
	2,4-D	(2,4-dichlorophenoxy)acetic acid
20	3,4-DA	(3,4-dichlorophenoxy)acetic acid
	dalapon	2,2-dichloropropanoic acid
	dazomet	tetrahydro-3,5-dimethyl-2H-1,3,5-thiadiazine-2-thione
	2,4-DB	4-(2,4-dichlorophenoxy)butanoic acid
	3,4-DB	4-(3,4-dichlorophenoxy)butanoic acid
25	DCB	1,2-dichlorobenzene
	DCPA	dimethyl 2,3,5,6-tetrachloro-1,4-benzenedicarboxylate
	DCU	N,N'-bis(2,2,2-trichloro-1-hydroxyethyl)urea
	2,4-DEB	2-(2,4-dichlorophenoxy)ethyl benzoate

	delachlor	2-chloro-N-(2,6-dimethylphenyl)-N-[(2-methylpropoxy)methyl]acetamide
	2,4-DEP	tris[2-(2,4-dichlorophenoxy)ethyl]phosphite
	desmedipham	ethyl[3-[[[(phenylamino)carbonyl]oxy]phenyl]carbamate
	desmetryn	N-methyl-N'-(1-methylethyl)-6-(methylthio)-1,3,5-triazine-2,4-diamine
5	diallate	S-(2,3-dichloro-2-propenyl) bis(1-methylethyl)carbamothioate
	dicamba	3,6-dichloro-2-methoxybenzoic acid
	dichlobenil	2,6-dichlorobenzonitrile
	dichlormate	3,4-dichloro benzenemethanol methylcarbamate
	dichlorprop	(±)-2-(2,4-dichlorophenoxy)propanoic acid
10	diclofop	(±)-2-[4-(2,4-dichlorophenoxy)phenoxy]propanoic acid
	dicryl	N-(3,4-dichlorophenyl)-2-methyl-2-propenamide
	diethatyl	N-(chloroacetyl)-N-(2,6-diethylphenyl)glycine
	diclosulam	N-(2,6-dichlorophenyl)-5-ethoxy-7-fluoro[1,2,4]triazolo[1,5-c]pyrimidine-2-sulfonamide
	difenopenten	(E)-(±)-4-[4-[4-(trifluoromethyl)phenoxy]phenoxy]-2-pentenoic acid
15	difenoxuron	N'-[4-(4-methoxyphenoxy)phenyl]-N,N-dimethylurea
	difenzoquat	1,2-dimethyl-3,5-diphenyl-1H-pyrazolium
	dimethachlor	2-chloro-N-(2,6-dimethylphenyl)-N-(2-methoxyethyl)acetamide
	dimethametryn	N-(1,2-dimethylpropyl)-N'-ethyl-6-(methylthio)-1,3,5-triazine-2,4-diamine
	dinitramine	N3,N3-diethyl-2,4-dinitro-6-(trifluoromethyl)-1,3-benzenediamine
20	dinosam	2-(1-methylbutyl)-4,6-dinitrophenol
	dinoseb	2-(1-methylpropyl)-4,6-dinitrophenol
	dinoterb	2-(1,1-dimethylethyl)-4,6-dinitrophenol
	diphenamid	N,N-dimethyl-α-phenyl benzeneacetamide
	dipropetryn	6-(ethylthio)-N,N'-bis(1-methylethyl)-1,3,5-triazine-2,4-diamine
25	diquat	6,7-dihydrodipyrido[1,2-a:2',1'-c]pyrazinediium ion
	dithiopyr	S,S-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate
	diuron	N'-(3,4-dichlorophenyl)-N,N-dimethylurea
	DNOC	2-methyl-4,6-dinitrophenol

	3,4-DP	2-(3,4-dichlorophenoxy) propanoic acid
	DSMA	disodium salt of MAA
	EBEP	ethyl bis (2-ethylhexyl)phosphinate
	eglinazine	N-(4-chloro-6-ethylamino-1,3,5-triazin-2-yl)glycine
5	endothall	7-oxabicyclo[2.2.1]heptane-2,3-dicarboxylic acid
	endothal-sodium	Sodium salt of endothal
	EPTC	S-ethyl dipropyl carbamothioate
	erbon	2-(2,4,5-trichlorophenoxy)ethyl-2,2-dichloropropanoate
	ethalfuralin	N-ethyl-N-(2-methyl-2-propenyl)-2,6-dinitro-4-(trifluoromethyl)benzenamine
10	ethametsulfuron	2-[[[[[4-ethoxy-6-(methylamino)-1,3,5-triazin-2-yl]amino]carbonyl]amino]sulfonyl]benzoic acid
	ethidimuron	N-(5-ethylsulfonyl-1,3,4-thiadiazol-2-yl)-N,N'-dimethylurea
	ethiolate	S-ethyl diethylcarbamothioate
	ethofumesate	(±)-2-ethoxy-2,3-dihydro-3,3-dimethyl-5-benzofuranyl methanesulfonate
	EXD	diethyl thioperoxydicarbonate
15	fenac	2,3,6-trichlorobenzeneacetic acid
	fenoxaprop	(±)-2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoic acid
	fenuron	N,N-dimethyl-N'-phenylurea
	fenuron TCA	salt of fenuron and TCA
	flamprop	N-benzoyl-N-(3-chloro-4-fluorophenyl)-DL-alanine
20	fluazifop	(±)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid
	fluazifop-P	(R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid
	fluchloralin	N-(2-chloroethyl)-2,6-dinitro-N-propyl-4-(trifluoromethyl)benzenamine
	flumetsulam	N-(2,6-difluorophenyl)-5-methyl[1,2,4]triazolo[1,5-a]pyrimidine-2-sulfonamide
	flumiclorac	[2-chloro-4-fluoro-5(1,3,4,5,6,7-hexahydro-1,3-dioxo-2H-isoindol-2-yl)phenoxy]acetic acid
25	flumioxazin	2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione
	fluometuron	N,N-dimethyl-N'-[3-(trifluoromethyl)phenyl]urea

	fluorochloridone	3-chloro-4-(chloromethyl)-1-[3-(trifluoromethyl)phenyl]-2-pyrrolidinone
	fluorodifen	2-nitro-1-(4-nitrophenoxy)-4-trifluoromethylbenzene
	fluoroglycofen	carboxymethyl 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate
	flupropacil	1-methylethyl 2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]benzoate
5	flupyrsulfuron	2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-6-(trifluoromethyl)-3-pyridinecarboxylic acid
	fluridone	1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone
	fluroxypyr	[(4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic acid
	flurtamone	(±)5(methylamino)-2-phenyl-4-[3-(trifluoromethyl)phenyl]-3(2H)-furanone
	fomesafen	5-[2-chloro-4-(trifluoromethyl)phenoxy]-N-(methylsulfonyl)-2-nitrobenzamide
10	fosamine	ethyl hydrogen (aminocarbonyl)phosphonate
	glufosinate	2-amino-4-(hydroxymethylphosphinyl)butanoic acid
	glyphosate	N-(phosphonomethyl)glycine
	halosafen	5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-N-(ethylsulfonyl)-2-nitrobenzamide
	haloxyfop	(±)-2-[4-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid
15	hexaflurate	potassium hexafluoroarsenate
	hexazinone	3-cyclohexyl-6(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H)-dione
	imazamethabenz	(±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-4(and 5)-methylbenzoic acid (3:2)
	imazamox	2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid
	imazapyr	(±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid
20	imazaquin	2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid
	imazethapyr	2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid
	ioxynil	4-hydroxy-3,5-diiodobenzonitrile

	ipazine	6-chloro-N,N-diethyl-N'-(1-methylethyl)-1,3,5-triazine-2,4-diamine
	IPX	O-(1-methylethyl)carbonodithioate
	isocarbamid	N-(2-methylpropyl)-2-oxo-1-imidazolidinecarboxamide
	isocil	5-bromo-6-methyl-3-(1-methylethyl)-2,4(1H,3H)-pyrimidinedione
5	isomethiozin	6-(1,1-dimethylethyl)-4-[(2-methylpropylidene)amino]-3-(methylthio)-1,2,4-triazin-5-(4H)-one
	isopropalin	4-(1-methylethyl)-2,6-dinitro-N,N-dipropylbenzenamine
	isoproturon	N,N-dimethyl-N'-[4-(1-methylethyl)phenyl]urea
	isouron	N'-[5-(1,1-dimethylethyl)-3-isoxazolyl]-N,N-dimethylurea
	isoxaben	N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide
10	karbutilate	3-[[[(dimethylamino)carbonyl]amino]phenyl (1,1-dimethylethyl)carbamate
	KOCN	potassium cyanate
	lactofen	(±)-2-ethoxy-1-methyl-2-oxoethyl 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate
	lenacil	3-cyclohexyl-6,7-dihydro-1H-cyclopentapyrimidine-2,4(3H,5H)-dione
	linuron	N'-(3,4-dichlorophenyl)-N-methoxy-N-methylurea
15	MAA	methylarsonic acid
	MAMA	monoammonium salt of MAA
	maleic hydrazide	1,2-dihydro-3,6-pyridazinedione
	MCPA	(4-chloro-2-methylphenoxy)acetic acid
	MCPB	4-(4-chloro-2-methylphenoxy)butanoic acid
20	mecoprop	(±)-2-(4-chloro-2-methylphenoxy)propanoic acid
	mefluidide	N-[2,4-dimethyl-5-[(trifluoromethyl)sulfonyl]amino]phenyl]acetamide
	metam-sodium	Sodium salt of metham
	metamitron	4-amino-3-methyl-6-phenyl-1,2,4-triazin-5(4H)-one
	methalpropalin	N-(2-methyl-2-propenyl)-2,6-dinitro-N-propyl-4-(trifluoromethyl)benzenamine
25	metham	methylcarbamodithioic acid
	methazole	2-(3,4-dichlorophenyl)-4-methyl-1,2,4-oxadiazolidine-3,5-dione
	methibenzuron	N-(2-benzothiazolyl)-N,N'-dimethylurea N-(3-methoxypropyl)-N'-(1-methylethyl)-6-(methylthio)-methoprotryn 1,3,5-triazine-2,4-diamine

	methyl bromide	bromomethane
	metobromuron	N'-(4-bromophenyl)-N-methoxy-N-methylurea
	metolachlor	(2-methoxy-1-methylethyl)acetamide 2-chloro-N-(2-ethyl-6-methylphenyl)-N-
	metosulam	N-(2,6-dichloro-3-methylphenyl)-5,7-dimethoxy[1,2,4]triazolo [1,5-a]pyrimidine-2-sulfonamide
5	metoxuron	N'-(3-chloro-4-methoxyphenyl)-N,N-dimethyl urea
	metribuzin	4-amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one
	metsulfuron	2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino] carbonyl]amino]sulfonyl]benzoic acid
	molinate	S-ethyl hexahydro-1H-azepine-1-carbothioate
	monalide	N-(4-chlorophenyl)-2,2-dimethylpentanamide
10	monolinuron	N'-(4-chlorophenyl)-N-methoxy-N-methylurea
	monuron	N'-(4-chlorophenyl)-N,N-dimethylurea
	monuron TCA	salt of monuron and TCA
	MSMA	monosodium salt of MAA
	napropamide	N,N-diethyl-2-(1-naphthalenyloxy)propanamide
15	naptalam	2-[(1-naphthalenylamino)carbonyl]benzoic acid
	neburon	N-butyl-N'-(3,4-dichlorophenyl)-N-methylurea
	nicosulfuron	2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]- N,N-dimethyl-3-pyridinecarboxamide
	nitralin	4-(methylsulfonyl)-2,6-dinitro-N,N-dipropylbenzenamine
	nitrofen	2,4-dichloro-1-(4-nitrophenoxy)benzene
20	nitrofluorfen	2-chloro-1-(4-nitrophenoxy)-4-(trifluoromethyl)benzene
	norea	N,N-dimethyl-N'-(octahydro-4,7-methano-1H-inden-5-yl)urea 3aa,4a,5a,7a,7aa-isomer
	norflurazon	4-chloro-5-(methylamino)-2-(3-(trifluoromethyl)phenyl)-3(2H)- pyridazinone
	OCH	2,3,4,4,5,5,6,6-octachloro-2-cyclohexen-1-one
	oryzalin	4-(dipropylamino)-3,5-dinitrobenzenesulfonamide
25	oxadiazon	3-[2,4-dichloro-5-(1-methylethoxy)phenyl]-5-(1,1-dimethylethyl)- 1,3,4-oxadiazol-2-(3H)-one
	oxyfluorfen	2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-(trifluoromethyl)benzene

	paraquat	1,1'-dimethyl-4,4'-bipyridinium ion
	PBA	chlorinated benzoic acid
	PCP	pentachlorophenol
	pebulate	S-propyl butylethylcarbamothioate
5	pelargonic acid	nonanoic acid
	pendimethalin	N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine
	perfluidone	1,1,1-trifluoro-N[2-methyl-4-(phenylsulfonyl)phenyl]methanesulfonamide
	phenisopham	3-[[[(1-methylethoxy)carbonyl]amino]phenyl ethylphenylcarbamate
	phenmedipham	3-[(methoxycarbonyl)amino]phenyl (3-methylphenyl)carbamate
10	picloram	4-amino-3,5,6-trichloro-2-pyridinecarboxylic acid
	piperophos	S-[2-(2-methyl-1-piperidiny)-2-oxoethyl]O,O-dipropyl phosphorodithioate
	PMA	(acetato-O)phenylmercury
	potassium azide	potassium azide
	primisulfuron	2-[[[[[4,6-bis(difluoromethoxy)-2-pyrimidinyl]amino]carbonyl]amino]sulfonyl]benzoic acid
15	procyazine	2-[[[4-chloro-6-(cyclopropylamino)-1,3,5-triazine-2-yl]amino]-2-methylpropanenitrile
	prodiamine	2,4 dinitro-N3,N3-dipropyl-6-(trifluoromethyl)-1,3-benzenediamine
	profluralin	N-(cyclopropylmethyl)-2,6-dinitro-N-propyl-4-(trifluoromethyl)benzenamine
	proglinazine	N-[4-chloro-6-(1-methylethylamino)-1,3,5-triazine-2-yl]glycine
	prometon	6-methoxy-N,N'-bis(1-methylethyl)-1,3,5-triazine-2,4-diamine
20	prometryn	N,N'-bis(1-methylethyl)-6-(methylthio)-1,3,5-triazine-2,4-diamine
	pronamide	3,5-dichloro (N-1,1-dimethyl-2-propynyl)benzamide
	propachlor	2-chloro-N-(1-methylethyl)-N-phenylacetamide
	propanil	N-(3,4-dichlorophenyl)propanamide
	propaquizafop	(R)-2-[[[(1-methylethylidene)amino]oxy]ethyl 2-[4-[(6-chloro-2-quinoxalinyloxy]phenoxy]propanoate
25	propazine	6-chloro-N,N'-bis(1-methylethyl)-1,3,5-triazine-2,4-diamine
	propham	1-methylethyl phenylcarbamate
	prosulfalin	N-[[[4-(dipropylamino)-3,5-dinitrophenyl]sulfonyl]-S,S-dimethylsulfilimine

	proxan-sodium	sodium salt of IPX
	prynachlor	2-chloro-N-(1-methyl-2-propynyl)-N-phenylacetamide
	pyrazon	5-amino-4-chloro-2-phenyl-3(2H)-pyridazinone
	pyriclor	2,3,5-trichloro-4-pyridinol
5	pyridate	O-(6-chloro-3-phenyl-4-pyridazinyl) S-octyl carbonothioate
	pyrithiobac	2-chloro-6-[(4,6-dimethoxy-2-pyrimidinyl)thio]benzoic acid
	quinclorac	3,7-dichloro-8-quinolinecarboxylic acid
	quinonamid	2,2-dichloro-N-(3-chloro-1,4-dihydro-1,4-dioxo-2-naphthalenyl)acetamide
	quizalofop	(±)-2-[4-[(6-chloro-2-quinoxalinyloxy]phenoxy]propanoic acid
10	rimsulfuron	N-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-3-(ethylsulfonyl)-2-pyridinesulfonamide
	sebumeton	N-ethyl-6-methoxy-N'-(1-methylpropyl)-1,3,5-triazine-2,4-diamine
	sethoxydim	2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one
	sesone	2-(2,4-dichlorophenoxy)ethyl hydrogen sulfate
	siduron	N-(2-methylcyclohexyl)-N'-phenylurea
15	silvex	2-(2,4,5-trichlorophenoxy)propanoic acid
	simazine	6-chloro-N,N'-diethyl-1,3,5-triazine-2,4-diamine
	simeton	N,N'-diethyl-6-methoxy-1,3,5-triazine-2,4-diamine
	simetryn	N,N'-diethyl-6-(methylthio)-1,3,5-triazine-2,4-diamine
	sodium arsenite	sodium arsenite
20	sodium azide	sodium azide
	sodium chlorate	sodium chlorate
	solan	N-(3-chloro-4-methylphenyl)-2-methylpentanamide
	sulfentrazone	N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide
	sulfometuron	2-[[[(4,6-dimethyl-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]benzoic acid
25	swep	methyl(3,4-dichlorophenyl)carbamate
	2,4,5-T	(2,4,5-trichlorophenoxy)acetic acid
	2,4,5-TB	4-(2,4,5-trichlorophenoxy)butanoic acid

	2,3,6-TBA	2,3,6-trichlorobenzoic acid
	TCA	trichloroacetic acid
	tebuthiuron	N-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-N,N'-dimethylurea
	terbacil	5-chloro-3-(1,1-dimethylethyl)-6-methyl-2,4(1H,3H)-pyrimidinedione
5	terbuchlor	N-(butoxymethyl)-2-chloro-N-[2-(1,1-dimethylethyl)-6-methylphenyl]acetamide
	terbumeton	N-(1,1-dimethylethyl)-N'-ethyl-6-methoxy-1,3,5-triazine-2,4-diamine
	terbuthylazine	6-chloro-N-(1,1-dimethylethyl)-N'-ethyl-1,3,5-triazine-2,4-diamine
	terbutol	2,6-bis(1,1-dimethylethyl)-4-methylphenyl methylcarbamate
	terbutryn	N-(1,1-dimethylethyl)-N'-ethyl-6-(methylthio)-1,3,5-triazine-2,4-diamine
10	tetrafluron	N,N-dimethyl-N'-[3-(1,1,2,2-tetrafluoroethoxy)phenyl]urea
	thiazafluron	N,N'-dimethyl-N-[5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl]urea
	thiazopyr	methyl-2-(difluoromethyl)-5-(4,5-dihydro-2-thiazolyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3-pyridinecarboxylate
	thifensulfuron	3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylic acid
	thiobencarb	S-[(4-chlorophenyl)methyl]diethylcarbamothioate
15	2,2,3-TPA	2,2,3-trichloropropionic acid
	triallate	S-(2,3,3-trichloro-2-propenyl) bis(1-methylethyl)carbamothioate
	triasulfuron	2-(2-chloroethoxy)-N-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl] benzenesulfonamide
	tribenuron	2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino]sulfonyl]benzoic acid
	tricamba	2,3,5-trichloro-6-methoxy benzoic acid
20	triclopyr	[(3,5,6-trichloro-2-pyridinyl)oxy]acetic acid
	tridiphane	2-(3,5-dichlorophenyl)-2-(2,2,2-trichloroethyl)oxirane
	trietazine	6-chloro-N,N,N'-triethyl-1,3,5-triazine-2,4-diamine
	trifluralin	2,6-dinitro-N,N-dipropyl-4-(trifluoromethyl)benzenamine
	triflusulfuron	2-[[[(4-(dimethylamino)-6-(2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-3-methylbenzoic acid
25	trimeturon	methyl N'-(4-chlorophenyl)-N,N-dimethylcarbamidate
	tritac	1-[(2,3,6-trichlorophenyl)methoxy]-2-propanol

vernolate	S-propyl dipropylcarbamothioate
xylachlor	2-chloro-N-(2,3-dimethylphenyl)-N-(1-methylethyl)acetamide

*source <http://piked2.agn.uiuc.edu/wssa/subpages/herbicide/herbtab.htm>

Also within the scope of the present invention is the stimulation of the activity of an ecto-phosphatase and an ABC transporter by the over-expression of a regulatory molecule which may act by up-regulating the expression levels or by post-translationally modifying the ecto-phosphatase and the ABC transporter. Such activating regulatory molecules (e.g. calmodulin) may be over-expressed alone or together with the over-expression of the ecto-apyrase and the ABC transporter or any other combination.

Particular embodiments of the invention include polynucleotides that encode MDR-ABC transporter polypeptides, ecto-phosphatase polypeptides, and stimulatory regulatory polypeptides which are capable of stimulating the efflux of drug molecules from the cells, thus conferring drug resistance. The term polynucleotide encompasses nucleic acid molecules that encode a complete protein, as well as nucleic acid molecules that encode peptides, polypeptides, or fragments of a complete protein. The polynucleotides may comprise the wild-type allele (or a portion of such an allele) of a functional peptide ABC transporter and ecto-phosphatase, or they may comprise a mutated allele of such genes. The preferred polynucleotides encode the wild-type plant, *Arabidopsis thaliana*, AtPGP-1 ABC transporter (GenBank accession # X61370); wild-type *Homo sapiens* Pgp ABC transporter (GenBank accession # M29432); wild-type *Homo sapiens* MRP- β ABC transporter (PCT WO 98/46736); wild-type yeast, *Saccharomyces cerevisiae*, transporter STS1 (GenBank accession # X75916); wild-type yeast, *Saccharomyces cerevisiae*, transporter Pdr5p (GenBank accession # 1420383); wild-type *Aspergillus fumigatus* Afu-MDR1 ABC transporter (U.S. Patent No. 5,705,352); wild-type bacterial, *Lactococcus lactis*, transporter LmrA (GenBank accession # U63741); wild-type plant, *Pisum sativum*, ecto-phosphatase, apyrase (GenBank accession # Z32743); and for wild-type *Homo sapiens* apyrase (GenBank accession # AF034840); other ecto-phosphatases include *Homo sapiens* CD39L2 (GenBank accession # AF039916); *Homo sapiens* CD39L3 (GenBank accession # AF039917); *Homo sapiens* CD39L4 (GenBank accession # AF039918); and *Homo sapiens* ATP diphosphohydrolase (GenBank accession # HSU87967).

In one embodiment of the invention, the polynucleotides are operably linked to regulatory sequences sufficient to permit the expression of the polynucleotide in a host cell. Such polynucleotides may be incorporated into nucleic acid vectors that are sufficient to permit either the propagation or maintenance of the polynucleotide within a host cell, and
5 expression therein. The nature of the regulatory elements will depend upon the host cell, and the desired manner of expressing the polynucleotides.

The invention particularly contemplates providing the polynucleotides to plants. Suitable plants include, but are not limited to, species from the genera *Fragaria*,
Lotus, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*, *Geranium*,
10 *Manihot*, *Daucus*, *Arabidopsis*, *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Datura*,
Hyoscyamus, *Lycopersicon*, *Nicotiana*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*,
Antirrhinum, *Hemerocallis*, *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ranunculus*,
Senecio, *Salpiglossis*, *Cucumis*, *Bromelia*, *Glycine*, *Lolium*, *Zea*, *Triticum*, *Sorghum*,
Ipomoea, *Passiflora*, *Cyclamen*, *Malus*, *Prunus*, *Rosa*, *Rubus*, *Populus*, *Santalum*, *Allium*,
15 *Lilium*, *Narcissus*, *Ananas*, *Arachis*, *Phaseolus*, *Pisum*, *Oryza*, *Hordeum*, *Gossypium*.

Preferred prokaryotic vectors for subcloning and production of DNA include plasmids such as those capable of replication in *E. coli* such as, for example, pBR322, ColE1, psC101, pACYC184, such as those disclosed by Maniatis, T., et al. (In: *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y.
20 (1982)); pET11a, pET3a, pET11d, pET3d, pET22d, pET12a, pET28a, and other pET variants (Novagen); pCDNA3, pCDNA1 (InVitrogen).

A variety of methods may be used to introduce the polynucleotides of the present invention into a plant cell. Some examples include, but are not limited to, microinjection directly into the plant embryo cells or introduced by electroporation as
25 described in Fromm et al., 1985, *Proc. Natl. Acad. Sci. USA* 82:5824-5828; direct precipitation using polyethylene glycol as described in Paszkowski et al., 1984, *EMBO J.* 3:2717-2722; in the case of monocotyledonous plants, transformation of pollen with total DNA or an appropriate functional clone and the pollen can then be used to produce progeny by sexual reproduction; introduction of polynucleotides with the Ti plasmid of *Agrobacterium*
30 *tumefaciens* which provides a means for introducing DNA into plant cells (Horsch et al., 1988, *Current Communications in Molecular Biology*, Cold Spring Harbor Press, Cold

Spring Harbor, N.Y., pp 13-19); introduction of polynucleotides with the cauliflower mosaic virus (CaMV) (U.S. Patent No. 4,407,956).

5 A particularly useful Ti plasmid-based vector is Pkylx71. Schardl, C. et al., 1987, *Gene* 61:1-11. This vector utilizes the natural transfer properties of the Ti plasmid. A cloning vehicle such as pKYLX71 allows the insertion of a polynucleotide sequence into the expression cassette by a single recombination event.

The introduction of the transferred DNA (T-DNA) of the plasmid is accomplished by infecting root calli from Ws ecotype *Arabidopsis thaliana* with *Agrobacterium tumefaciens* under kanamycin selection. The calli are then developed further
10 into plants. Valvekens, D., 1992, *Proc. Natl. Acad. Sci. USA* 85:5536-5540. Alternatively, shoot explants may be infected with the *Agrobacterium tumefaciens* bacteria. Under appropriate conditions, a ring of calli forms around the cut surface which is then transferred to growth medium, allowed to form shoots, roots and develop further into plants. Hooykass, P.J. J. et al., In: *Molecular Form and Function of the Plant Genome*, Plenum Press, N.Y. pp
15 655-667 (1984). Another alternative is to produce transformed plants using free DNA delivery. All plants from which protoplasts can be isolated and cultured to give whole regenerated plants can be transformed by the present invention so that whole plants are recovered which contain the introduced polynucleotide. Methods for generating plants from cultured protoplasts are described by Binding, H. In: *Plant Protoplasts*, CRC Press, Boca
20 Raton, pp. 21-37 (1985), incorporated herein by reference.

Efficient plant promoters that may be used to over-express the ABC transporters and the ecto-phosphatases include over-producing plant promoters such as the small subunit (ss) of the ribulose 1, 5 biphosphate carboxylase from soybean (Berry-Lowe et al., 1982, *J Molec. App. Gen.* 1:483-498), the promoter of the chlorophyll a/b binding protein,
25 and the CaMV promoter.

Parts obtained from the recombinant plant such as flowers, seeds, leaves, branches, bark, fruit, etc, are covered by the invention. Progeny, variants, and mutants of the recombinant plants are also included within the scope of this invention.

Conference of Drug Resistance in Microorganisms

The present invention is also directed to a method for the conference of drug resistance to microorganisms, including yeast and bacteria in part through the manipulation of the ATP gradient across biological membranes. In yeast and bacteria, the manipulation of extracellular ATP levels and the ATP gradient across biological membranes by the over-expression of a MDR-ABC transporter and/or an ecto-phosphatase may result in resistance to certain drugs. Such resistance is useful for the growth of microorganisms for biotechnological applications, *e.g.*, those used in heterologous protein production.

It is particularly advantageous to be able to produce microorganisms which are resistant to a variety of drugs for large scale fermentation procedures where contamination by microorganisms from the environment may threaten a costly procedure. Additionally, the present invention is useful to create resistant microorganism strains in small scale fermentation processes, industrial applications, as well as in selection systems for the production of recombinant microorganisms for research applications. Research applications may include the use of resistant microorganism strains to study alternative pathways, other than antibiotics, antifungal reagents, or other commonly used drugs which could effectively inhibit the growth of microorganisms involved in disease states of humans and animals.

In yeast, a system which could confer drug resistance may be preferred to current research techniques which utilize yeast strains deficient for certain amino acid production pathways. These deficient yeast are used to introduce foreign nucleic acids of interest having a nucleotide sequence encoding a protein or proteins capable of resurrecting a deficient amino acid production pathway. Selection occurs when the yeast is grown in media deficient in that particular amino acid. This method of conferring resistance to yeast may be costly, however, since this requires that the yeast be grown in expensive cocktails of the amino acids in which they are deficient. In certain embodiments of the present invention, a cloning system in yeast confers drug resistance to the yeast coupled to the introduction of a nucleic acid molecule of interest. Such resistance may be constitutive or inducible. The yeast may then be selected by the introduction of inexpensive drugs to which the recombinant yeast would be resistant.

In other embodiments of the invention, bacteria may be produced with increased resistance to certain drugs in order to facilitate the production and to provide a system which allows for selection of bacteria based on another mechanism other than

antibiotic resistance. Such resistance may be constitutive or inducible and may be particularly useful in large scale fermentation where contamination by other microorganisms is more likely to occur.

Also contemplated by the present invention is the development of
5 microorganisms which grow in soil (soil flora), particularly those designed to interact with herbicide resistant plants. The soil flora may be engineered with the same resistance to toxins as the plants with which they are engineered to react.

Additionally, the invention is directed to the development of
microorganisms which are resistant to multiple toxins (two-stage resistant microorganisms or
10 multiple-stage resistant microorganisms). The toxins could be presented to such two-stage resistant organisms or multiple-stage microorganisms simultaneously or at independent times. The present invention also contemplates the development of two-stage or multiple-stage resistant plants.

In one embodiment of the invention, the over-expression of an ecto-
15 phosphatase confers drug resistance in wild-type or genetically engineered microorganisms. This effect was seen in yeast cells over-expressing plant apyrase grown in the presence of cycloheximide, a potent inhibitor of protein expression.

In another embodiment of the invention, the over-expression of an ABC
transporter confers drug resistance in wild-type and genetically engineered microorganisms.
20 In a preferred embodiment, the ABC transporter which is over-expressed is the *Arabidopsis thaliana* ABC transporter AtPGP-1. This ABC transporter was able to confer resistance to yeast cells grown in the presence of cycloheximide.

In a further embodiment of the invention the affect of over-expression of
both an MDR-ABC transporter and an ecto-phosphatase is to enhance the ATP gradient
25 across biological membranes and thus stimulate the resistance to certain antimicrobial agents. In a particularly preferred embodiment of the invention the MDR-ABC transporter which is over-expressed is the *Arabidopsis thaliana* AtPGP-1 and the ecto-phosphatase that is over-expressed is *Pisum sativum* apyrase.

The invention particularly contemplates, but is not limited to, the
30 conference of resistance in microorganisms to cycloheximide, antibiotics, antifungal agents, pheromones, heavy metals, flourescent dyes, DNA intercalating agents, products of plant

secondary metabolism such as polyphenolics and alkaloids, plant growth substances with antimicrobial properties, and the chemicals listed in Table 1 above.

In one embodiment of the invention, the nucleic acids are operably linked to regulatory sequences sufficient to permit the transcription of the nucleic acid in the microorganism of interest. Such constructs may be incorporated into nucleic acid vectors that are sufficient to permit either the propagation or maintenance of the nucleic acid and expression thereof within the host cell. The nature of the regulatory elements is dependent upon the host cell, and the desired manner of expressing the nucleic acid (e.g. constitutively or inducibly).

The invention particularly contemplates providing the nucleic acids of interest to bacteria and yeast. Suitable bacteria include both archaeobacteria, which are found in incommensurable environments such as bogs, ocean depths, salt brines, and hot acid springs (e.g. sulfur bacteria, extreme halophiles, methanogens), and eubacteria, which are the commonly encountered forms that inhabit soil, water, and larger living organisms (e.g. gram positive, anaerobic, blue-green algae, gram negative, and spirochetes). In a preferred embodiment, the bacteria are *Escherichia coli*. Suitable yeast include a large group of disparate organisms. Preferred species include the budding yeast, *Saccharomyces cerevisiae*, and the fission yeast, *Schizosaccharomyces pombe*.

Preferred prokaryotic vectors include, but are not limited to, plasmids such as those capable of replication in *E. coli*, for example, pBR322, ColE1, pSC101, pACYC 184 such as those disclosed by Maniatis, T., et al. (In: *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1982)); pET11a, pET3a, pET11d, pET3d, pET22d, pET12a, pET28a, and other pET variants (Novagen); pCDNA3, pCDNA1 (Invitrogen); pRR54, pRS303, pEGFP-1, pBluescript SK, pTrc99A,B,C and their derivatives (In: *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., Mass., USA (1998)); pGEX variants (Pharmacia) and bacteriophages (e.g. Lambda phages).

Preferred yeast vectors include plasmids such as those capable of replication in either *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*. These vectors include, but are not limited to, pYES2, pVT101, Yip5, Prp7, Yrp17, Pep13, Yep24, Ycp19, Ycp50, Ylp21, pYAC3, 2 μ m, pLG670. In: *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., Mass., USA (1998).

A variety of methods may be used to introduce the polynucleotide sequences into a microorganism. In bacteria for example, techniques such as transformation of plasmid DNA using calcium chloride competent cells, high efficiency competent cells, electroporation, or infection by bacteriophages as described in *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., Mass., USA (1998) may be used.

In yeast, methods to introduce polynucleotides can include, but are not limited to, the introduction of polynucleotides by integrative transformation, transformation by electroporation, spheroplast transformation, transformation using lithium acetate as described in *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., Mass., USA (1998) and PEG lithium acetate transformation procedure (Eble, R., 1992, *Biotechniques* 13:18-20).

Also within the scope of the present invention is the conference of drug resistance to eukaryotic cell lines grown in tissue culture, including insect cell lines and mammalian cell lines. The conference of drug resistance to eukaryotic cell lines may be useful in the use of such cell lines for the production of recombinant proteins, the study of chemotherapeutic resistance in cells from various sources, and in the study of toxic levels of drugs in certain resistant cell lines.

Preferred eukaryotic vectors include but are not limited to, viral vectors, naked nucleic acids, plasmids, shuttle vectors, complexes of nucleic acids and other molecules, such as polycations (e.g. cationic lipids), including those described in *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., Mass., USA (1998) for introduction of heterologous DNA in mammalian cells and those described in *Baculovirus Expression Vectors*; a laboratory manual, Oxford University Press, New York., N.Y. (1994) for introduction of heterologous DNA in insect cells.

Inhibition of Drug Resistance in Chemotherapy

In an alternative embodiment of the invention, modulation of the ATP gradient, specifically the suppression of the gradient, may be achieved by inhibiting the activity of the ecto-phosphatase and the ABC transporter. Suitable inhibitor mechanisms include, but are not limited to, the use of small molecules which may bind to and inhibit the activity of the ecto-phosphatase and small molecules which may bind to and inhibit the ABC transporter. Other suitable inhibitor mechanisms include, but are not limited to, the expression of anti-sense RNA molecules which may inhibit the transcription or translation of ecto-phosphatases and ABC transporters, as well as the expression of dominant negative mutants of the ecto-

phosphatase and the ABC transporters which may act to interfere with and inhibit the activity of their wild-type counterparts. Also within the scope of the invention is the over-expression of regulatory molecules which inhibit the activity of the ecto-phosphatase and the ABC transporter. The ecto-phosphatase may be inhibited alone or together with the ABC transporter.

Accordingly, it is one object of the present invention to provide molecules which down-regulate the activity of ecto-phosphatases either alone or together with ABC transporters.

The present invention provides for methods for the transcription of exogenous antisense RNA, *in vivo* or *in vitro*, comprising the administration of a polycistronic vector which may contain nucleic acid molecules from which may be transcribed an antisense RNA complementary to an ecto-phosphatase RNA molecule and an ABC transporter RNA molecule. The ecto-phosphatase nucleic acids and the ABC transporter nucleic acids may be operatively linked to a constitutive promoter or an inducible promoter (*e.g.* the constitutive major intermediate early promoter of cytomegalovirus or the inducible metallothionein promoter). Also within the scope of the invention are multiple vectors whereby the ecto-phosphatase nucleic acid and the ABC transporter nucleic acid are incorporated into separate vectors.

The present invention also provides for methods for the expression of exogenous regulatory molecules or small molecules, *in vivo* or *in vitro*, comprising the administration of a polycistronic vector or multiple vectors which may incorporate nucleic acid molecules encoding for regulatory proteins, or small molecules capable of inhibiting the activity of an ecto-phosphatase and an ABC transporter. These nucleic acids which encode for regulatory molecules may be operatively linked to either a constitutive promoter or an inducible promoter as described above.

Additionally, it is an object of the present invention to provide inhibitors of ecto-phosphatases (*e.g.* apyrase) in physiological compositions for modulating MDR states. Such physiological compositions comprise a small molecule capable of inhibiting an ecto-phosphatase and a physiologically acceptable carrier or diluent. As used herein, the term “physiologically acceptable carrier or diluent” means any and all solvents, dispersion media, antibacterial and antifungal agents, microcapsules, liposomes, cationic lipid carriers, isotonic and absorption delaying agents and the like which are not incompatible with the ecto-

phosphatase inhibitors. The use of such media and agents for physiologically active substances is well known in the art. Supplementary active ingredients may also be incorporated into the compositions.

5 In certain preferred embodiments, only the endogenous ecto-phosphatase is inhibited by antisense RNA, regulatory proteins, or small molecules. In other embodiments of the present invention, both the endogenous ecto-phosphatase and the ABC transporter are targeted for inhibition.

In particularly preferred embodiments, the ecto-phosphatase is human apyrase (e.g. GenBank accession # AF034840, AF039916, AF039917, AF039918,
10 HSU87967) and the MDR-ABC transporter is human MDR-1, (e.g. human P-glycoprotein MDR-1, GenBank accession # M29432 and human MRP- β PCT, publication WO 98/46736).

The nucleic acid molecules from which will be transcribed antisense RNA molecules described above or the nucleic acid molecules encoding for regulatory molecules as described above may be incorporated into any suitable cloning or expression vector, operably
15 linked to appropriate control elements (e.g. promoter elements, enhancer elements, ribosomal binding sites, polyadenylation sites, termination sites, etc.). Examples of such vectors include, but are not limited to, herpes simplex viral based vectors such as pHSV1 (Geller et al., 1990, *Proc. Natl. Acad. Sci. U.S.A.* 87:8950-8954); retroviral vectors such as MFG (Jaffee et al., 1993, *Cancer Res.* 53:2221-2226), and in particular Moloney retroviral vectors such as LN,
20 LNSX, LNCX, LXSX (Miller and Rosman, 1989, *Biotechniques* 7:980-989); vaccinia viral vectors such as MVA (Sutter and Moss, 1992, *Proc. Natl. Acad. Sci. U.S.A.* 89:10847-10851); adenovirus vectors such as pJM17 (Ali et al., 1994, *Gene Therapy* 1:367-384; Berker, 1988, *Biotechniques* 6:616-624; Wand and Finer, 1996, *Nature Medicine* 2:714-716); adeno-associated virus vectors such as AAV/neo (Mura-Cacho et al., 1992, *J. Immunother.* 11:231-
25 237); lentivirus vectors (Zufferey et al., 1997, *Nature Biotechnology* 15:871-875). Such vectors may be targeted to the tumor cells of interest as described in U.S. Patents 5,834,256, 5,843,742, 5,830,727, 5,814,500.

The inhibition of ecto-phosphatases either alone or together with the inhibition of ABC transporters is useful in the reduction of drug resistance in cells. In one
30 embodiment of the invention, the inhibition of ecto-phosphatases either alone or together with ABC transporters results in a loss of resistance to drug molecules used in chemotherapy. In

another embodiment of the invention, administration of such inhibitory molecules is in conjunction with the administration of chemotherapeutic agents in tumor cells.

Administration of the foregoing agents may be local or systemic, using a suitable physiological carrier. Other compounds which aid in the uptake or stability of these agents, or which have beneficial activity, may also be included in the formulations of the invention.

Inhibition of Drug Resistance in Microorganisms to Treat Infection

The present invention also relates to methods for inhibiting or ameliorating infection in animals and humans caused by microorganisms, particularly bacterial and fungal infections using inhibitory mechanisms against an ecto-phosphatase and an ABC transporter and modifying the ATP gradient across biological membranes. The invention is useful in the inhibition or amelioration of a wide range of infections including, but not limited to, gram-negative bacterial infection including gram-negative sepsis, gram-negative endotoxin-related hypotension and shock, rabies, cholera, tetanus, lymes disease, tuberculosis, *Candida albicans*, *Chlamydia*, etc. The invention is based, in part, on the unexpected result that when mutant yeast deficient in two potent extracellular ATP phosphatases were cultured in cycloheximide, they were not able to grow. Surprisingly, they were rescued by the over-expression of a plant MDR-ABC transporter AtPGP-1, suggesting that the inability to grow in the drug was caused by an inability to efflux the drug which was coupled to a deficiency in extracellular ATP phosphatase activity.

Drug sensitivity in microorganisms may be achieved by introducing nucleic acid molecules into bacteria and yeast (as described above) that are capable of conferring inhibition of the activity of an endogenous ecto-phosphatase and an ABC transporter. Such nucleic acid molecules may transcribe an antisense RNA complimentary to endogenous RNA for an ecto-phosphatase or an ABC transporter, encode for inhibitory regulatory proteins, or encode for inhibitory drug molecules. The inhibition or amelioration of the infections may involve the administration of an anti-microbial agent (such as an antibiotic or an antifungal agent) with the concurrent administration of the aforementioned nucleic acid molecules (which may be achieved through bacteriophages, etc). Additionally, inhibitors of ecto-phosphatases or ABC transporters may be administered via a physiologically acceptable carrier as described above.

Additionally, the present invention is useful in the development of genetic and epigenetic systems in humans for resistance to toxins from biological and non-biological sources. Such sources include, but are not restricted to, pathogens produced by microbial infections, pathogens and toxins derived from biological sources through human contrivance, environmental toxins not produced through biological action, and toxic substances created synthetically. In a particular embodiment, humans at risk for exposure would be vaccinated either with a gene therapy designed to bolster endogenous ATP gradients in human cells, or a chemical substance capable of enhancing the strength of the ATP gradient. In both instances, the target of the genetic or chemical therapy would be either the ABC transporter activity, ecto-phosphatase activity or both. In another embodiment of the invention, only the ABC transporter activity or the ecto-phosphatase activity in an infecting organism is diminished to inhibit drug efflux. Recombinant techniques may be used to introduce DNA sequences to the microorganism which encode for a small inhibitory molecule to either an ABC transporter or an ecto-phosphatase or both to cause the inhibition of drug efflux from the microorganism.

Ecto-phosphatase Inhibition


Since ecto-phosphatases have been shown by the present invention to be important actors in the modulation of the ATP gradient across biological membranes and thus useful in a variety of applications (e.g. the modulation of drug resistance), it is an object of the present invention to provide methods and assays for the identification of inhibitors of ecto-phosphatases (e.g. apyrase).

A high-throughput screen was developed to rapidly identify potential inhibitors for ecto-phosphatases and is described below in Example 6. This high-throughput screen is particularly useful, since no known specific inhibitors of the apyrase enzyme exist. Using the high throughput screen, ecto-phosphatase inhibitors are isolated by screening a small molecule library (e.g. a combinatorial library) for inhibitory activity to ecto-phosphatase (e.g. apyrase) activity. Once ecto-phosphatase inhibitory molecules are isolated from such a screen, the inhibitors may be further tested for their ability to specifically inhibit the ATPase activity of the ecto-phosphatase and to reduce drug resistance in cells.

The ecto-phosphatase inhibitory molecules of the present invention are chemically stable and physiologically active and include, *inter alia*, those molecules represented by Formulae I through XIX below.

I


5



10


II

15



III


20



25


IV

30




34

5




10



15

20




25

VII


VIII

30




35

5




10

15







20

25




30


36




XVI




XVII



XVIII



XIX



Preliminary pharmacophore studies revealed that the small molecules represented by Formulae I through XIX fall into five classes of compounds (sulfanamides, guanidines, aminothiazoles, thioketones and benzamides). Most of these chemical classes are found in other physiologically-active compounds, including those having pharmaceutical and therapeutic use. For example, sulfanimides are widely used as antibiotics. Additionally, studies for the isolation of small molecules capable of reversing MDR have described molecules belonging to two of the classes of molecules of the present invention (Medina et al., 1998, *Bioorg Med. Chem. Lett.* 8:2653-2656 and Dhamant et al., 1992, *J. Med. Chem.* 35:2481-2496). The molecules described by Medina et al. have been shown to affect MDR and the mode of action of the molecules is believed to involve tubulin interactions. The thiazine derivatives described by Dhamant et al. reverse the resistance in tumor cells to vincristine.

The ecto-phosphatase inhibitory molecules of the present invention are useful in reversing MDR in *Arabidopsis* plants and yeast. MDR reversal in plants and yeast cells may be shown by growing the cells in the presence of relevant drugs and in the presence and absence of the inhibitor. Cells which cannot grow in drug, in the presence of an ecto-phosphatase inhibitor, have a reversal in MDR. Additionally, the ecto-phosphatase inhibitory molecules of the present invention are useful in reversing drug resistance in mammalian cell lines (e.g. normal COS-7 cells and breast cancer tumor cells (e.g. HS5787, MB231 and MB435)) grown in the presence of a drug (e.g. a chemotherapeutic agent). MDR reversal in mammalian cells may be shown by using the fluorescent compound calcein-AM. Esterases present in cells cleave the aceto-methoxy ester (AM) from the calcein-AM and liberate calcein. Calcein is a fluorescent compound which is excitable by the 488 nm laser of a FACSCaliber flow cytometer (Becton Dickinson, Franklin Lakes, N.J.), while the uncleaved calcein-AM is not excitable. Wild type cells incubated in the presence of calcein-AM show a high level of fluorescence while MDR state cells, which efflux the calcein-AM faster than the cellular esterases can cleave it, do not show a high level of fluorescence. The mammalian cells can be tested for the reversal of MDR with the ecto-phosphatase inhibitors of the present invention by the amount of calcein fluorescence detected in the cells. Furthermore, the relative importance of the mammalian MDR gene and the mammalian apyrase gene in MDR can also be determined.

Specificity of the ecto-phosphatase inhibitors of the present invention may be tested with the screening assay described in Example 6 below. Inhibitors are tested for their ability to inhibit acid phosphatases, alkaline phosphatases, myosin phosphatases and the luciferase ATPase. The assays may be performed using techniques known in the art.

5 In one preferred embodiment, the ecto-phosphatase is an apyrase and the ecto-phosphatase inhibitor is a molecule selected from among molecules represented by the Formulae I through XIX. In another preferred embodiment, the ecto-phosphatase is apyrase and the ecto-phosphatase inhibitor is a molecule selected from among molecules represented by the Formulae I through V. In a particularly preferred embodiment, the ecto-phosphatase is
10 apyrase and the ecto-phosphatase inhibitor is a molecule selected from among molecules represented by Formula I and Formula II.

The ecto-phosphatase inhibitors of the present invention which are acidic or basic in nature can form a wide variety of salts with various inorganic and organic bases or acids, respectively. These salts may be physiologically acceptable for *in vivo* administration in plants
15 and animals, including humans. Salts of the acidic compounds of this invention are readily prepared by treating the acidic compound with an appropriate molar quantity of the chosen inorganic or organic base in an aqueous or suitable organic solvent and then evaporating the solvent to obtain the salt. Salts of the basic compounds of this invention can be obtained similarly by treatment with the desired inorganic or organic acid and subsequent solvent
20 evaporation and isolation. The skilled artisan can produce salts of the small molecules of the present invention using techniques known in the art.

The skilled artisan readily can determine the amount of the ecto-phosphatase inhibitor that is required to inhibit the ecto-phosphatase by measuring ATPase activity in the presence and absence of varying amounts of the inhibitor. Phosphatase activity can be determined by
25 assessing the dephosphorylation of ATP and liberation of phosphate as described below in Example 6. Additionally, parameters may be measured that are known to be associated with ecto-phosphatase activity to determine whether the molecule has ecto-phosphatase inhibitory activity. For example, ecto-phosphatase inhibitory activity may be measured in cells (*e.g.* plant, yeast, mammalian, tumor, etc. cell lines) by assessing the loss of resistance to drugs.
30 Furthermore, the ecto-phosphatase inhibitory molecules of the present invention may be tested for specific inhibitory activity to ecto-phosphatases versus general phosphatases or for specific inhibitory activity for a particular ecto-phosphatase activity (*e.g.* apyrase).

Additionally, as stated above, the ecto-phosphatase inhibitory molecules of the present invention are useful in reversing MDR. Such a reversal has several applications including reducing resistance to chemotherapeutic agents in tumor cells and reducing resistance to antimicrobial agents in microorganisms.

5 Inhibition of ecto-phosphatases is useful in industrial applications as well. For example, one of the most sensitive and cost effective ways of determining the titer of microbia in soil, sludge, blood, food, and textiles is the luciferase assay which allows for the estimation of microbial biomass through the determination of precise concentrations of ATP. The sensitivity of the assay requires that "background" ATP or nonmicrobial ATP present in the
10 system as a consequence of the source of the sample be separated from the ATP used in the microbe count. The removal of background ATP is accomplished using the ecto-phosphatase, apyrase. After removal of the background ATP with apyrase, the apyrase must be removed or inactivated. General techniques for removal could be improved and simplified with a method of inactivating the apyrase by adding a specific apyrase inhibitor of the present invention.

15 The present invention also provides physiologically acceptable compositions comprising an ecto-phosphatase inhibitor of the present invention and a physiologically acceptable carrier or diluent as described above. The use of such physiologically acceptable carriers or diluents are well known in the art. Formulation of such physiological compositions can be made using known procedures, e.g. according to Remington's Pharmaceutical
20 Sciences, 17th ed., Mack Publishing Co., Easton, Pa. Formulation of the compounds of the present invention may be stable under the conditions of manufacture and storage and must be preserved against contamination by microorganisms. Contamination can be avoided using antimicrobial (*e.g.* antibacterial and antifungal) agents.

 The physiological forms of the compounds of the invention suitable for administration
25 include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. Typical carriers include a solvent or dispersion medium containing, for example, water buffered aqueous solutions (*i.e.* biocompatible buffers), ethanol, polyols such as glycerol, propylene glycol, polyethylene glycol, suitable mixtures thereof, surfactants, and vegetable oils. Isotonic agents such as
30 sugars or sodium chloride may be incorporated into the subject compositions.

The present invention is further illustrated by the following examples which in no way should be construed as being further limiting. The contents of all references cited throughout this application are hereby expressly incorporated by reference.

5 EXAMPLE 1: OVER-EXPRESSION OF ECTO-PHOSPHATASE DOES NOT INCREASE
10 THE CELLULAR UPTAKE OF ADENOSINE

MATERIALS AND METHODS

Transgenic Plant Construction: psNTP9 (*Pisum Sativum* apyrase, GenBank accession #Z32743) was subcloned as a Sall to XbaI fragment into pKYLX71 (Schardl et al, 1987, *supra.*). This plasmid was transformed into *A. tumefaciens* GV3101 [pMP90] pKYLX71
10 (Koncz, C. and Shell, J., 1986, *Mol. Gen. Genet.* 204:383-396.), which was used to infect root calli from Ws ecotype *Arabidopsis thaliana* under kanamycin selection (Valvekens, D. et al., 1992, *Proc. Natl. Acad. Sci. USA* 85:5536-5540.). Four individual lines, obtained from separate calli, were propagated to the third generation (T3).
15 Subcellular Apyrase Distribution in Pea: Etiolated pea plumules served as the tissue source for nuclei and cytoplasm isolation as described by Chen and Roux (*Plant Physiol.* 81:609-612 (1986)). Plasma membrane was prepared from 30 g of pea root tissue (Zhu Mei Jun and Chen Jia, 1995, *Acta Botanica Sinica* 37:942-949). Western analysis was performed on 15-30 µg of protein from cytoplasm, plasma membrane and nuclei using a polyclonal anti-apyrase
20 antibody raised against the purified pea protein (Tong, C. et al., 1993, *Plant Physiol.* 101:1005-1011). To determine the orientation of the pea apyrase in the pea plasma membrane, outside-out vesicles were prepared (Short et al., *supra.*), and the accessibility of the enzyme was determined by selective trypsin proteolysis, or membrane shaving, followed by activity assays and western blotting.
25 Phosphate uptake experiments and growth assays: In all experiments the growth media did not contain sugar, and plants were grown in sterile culture at 22° C under 150-200 µE of continuous light. Unless otherwise noted, a standard 0.8% agar medium (Becton Dickenson, Cockeysville, Md.) containing 100 µM phosphate was used for uptake assays (Somerville, C. et al., 1982, *Methods in Chloroplast Biology*, Elsevier Biomedical Press, Amsterdam, pp 129-
30 138). Plants used for the phosphate uptake experiments were grown singly in 1 ml of the standard agar medium for 15 days prior to the experiment. On the day of the experiment, 10 µCi ³²P was applied to the side of the culture dish and allowed to diffuse through the agar.

The lids of 95 mm x 15 mm tissue culture dishes (Fisher, Pittsburgh, Pa.) were removed to facilitate transpiration. After 18 hours, the plants were removed from the medium. The aerial portions of the plant not in contact with the agar were weighed and counted by liquid scintillation. For each plant the entire root system was carefully pulled from the agar and washed in ice cold water prior to scintillation counting. To measure the transport of the products of ATP hydrolysis by the transgenic plants overexpressing apyrase and by wild-type plants, [2,8-³H]ATP, [α -³²P]ATP, and [γ -³²P] ATP (Amersham) were fed to 15-day-old plants in separate treatments. All treatments were analyzed for significance in a T-test ($n > 4-6$ for all groups, * $P < 0.05$, error bars = s.e.m.).

RESULTS

Detection of the pea apyrase in nuclei and in purified plasma membrane: By immunoblot assay, the pea apyrase was found to be associated with nuclei and with purified plasma membranes but not with the cytoplasm (Figure 1A). The contents of the lanes in Figure 1A are as follows: Lane 1, cytoplasm; Lane 2, purified plasma membrane; Lane 3, purified nuclei; and Lane 4, pre-immune control of nuclei. Protease treatment destroyed both apyrase activity and antigenicity in outside-out plasma membrane vesicles. After trypsin treatment, the exterior face of the vesicle showed 30% of the ecto-phosphatase activity of the untreated sample. Endo-phosphatase activities were retained after trypsin treatment, indicating that the digest occurred exclusively on the exterior face of the membrane. These data indicated that the ecto-apyrase was in fact being expressed in the extracellular matrix (ECM).

Enhanced Growth of Plants Over-Expressing Apyrase: Three of the four transgenic plant lines constitutively expressed psNTP9 under the control of the cauliflower mosaic virus 35S promoter and over an 18 hour period showed two to five times as much phosphate accumulation in shoots as wild type (Figure 1B); Top, the total phosphate accumulated in the shoots of three independent transformants in an 18 hour ³²P uptake assay at 2 mM phosphate; Bottom, a corresponding immunoblot performed on equal amounts of protein isolated from the ECM of three week-old wild-type *Arabidopsis thaliana* and the psNTP9 transgenics. Apyrase expressing plants also showed four times as much phosphatase activity in the extracellular matrix as the wild-type (Figure 1C). (Note, OE1 in the figure stands for over-expression 1 transgenic line).

Transgenic plants preferentially transport the gamma phosphate of ATP: In order to address whether over-expression of ecto-apyrase was stimulating the adenosine salvage pathway, the intracellular uptake of adenosine was measured both in the presence and absence of the over-expression of apyrase. The inability of apyrase to translocate either extracellular AMP or adenosine was demonstrated by the low level of radiolabel accumulated in the transgenic plants fed [2,8³H]ATP and [α ³²P]ATP (Figure 2). The complete dephosphorylation of [2,8³H]ATP would result in a radiolabelled adenosine molecule while the complete dephosphorylation of [α ³²P]ATP would result in a non-labeled adenosine label. Figure 2A illustrates that plants overexpressing apyrase did not translocate radiolabelled adenosine (or byproducts of the dephosphorylation of [2,8³H]ATP) any more efficiently than plants not overexpressing apyrase (wild-type plants). Figure 2B illustrates that plants overexpressing apyrase did not translocate AMP (or the byproducts of the dephosphorylated [α ³²P]ATP) any more efficiently than wild-type plants. In comparison, feeding experiments where the γ phosphate was labeled, the transgenics accumulated three times the amount of labeled phosphate as the wild-type (Figure 2C). These data show that the over-expression of apyrase does not induce an increase in the uptake of adenosine and therefore its over-expression does not act to stimulate the adenosine salvage pathway.

EXAMPLE 2: ECTO-PHOSPHATASE IS INVOLVED IN DRUG RESISTANCE IN

YEAST AND PLANTS

MATERIALS AND METHODS

Expression of AtPGP-1 in yeast: The AtPGP-1 cDNA (*Arabidopsis thaliana* MDR gene, accession #X61370) was subcloned into pVT101 downstream of the ADH promoter to create the AtPGP-1/pVT101 construct. AtPGP-1/pVT101 and pVT101 were transformed into *Saccharomyces cerevisiae* INVSC1 (genotype: *MAT α* , *his3- Δ 1*, *leu2*, *trp1-289*, *ura3-52*) and YMR4 (genotype: *MAT α his3-11,15*, *leu2-3*, *112ura3 Δ 5*, *can Res pho5*, *3::ura3 Δ 1*) by a PEG lithium acetate procedure (Eble, R., 1992, *Biotechniques* 13:18-20) and selected on uracil dropout medium.

Yeast Growth: Yeast were grown at 30° C under conditions of constant selection for uracil auxotrophy. YNB (Bio101, Vista, CA) supplemented with CSM (uracil dropout) and 2% glucose was used to grow strains having pVT101 constructs. Cycloheximide (Sigma Chemical, St. Louis, MO.) was added to liquid media or spread on solid media to achieve a

final concentration of 500 ng/ml. Nigericin (Sigma Chemical, St. Louis, MO.) was added to liquid media or spread on solid media to achieve a final concentration of 25 µg/ml. Yeast strains used in cycloheximide selection assays were always propagated in the presence of the cycloheximide on plates and then streaked onto new plates containing drug or no drug, such that induced resistance existed in each strain at the time of the start of the assay. For selection assays on plates, single colonies were streaked; for selection in liquid media 0.01 ml of saturated culture was added to fresh media containing the drug. The plates shown in figures were grown for 3-5 days before photographs were taken. Yeast selection assays in liquid media were quantitated by turbidity as measured by absorbance at OD₆₀₀.

Expression of apyrase and AtPGP-1 in plants: The expression of apyrase in plants is as described above in Example 1. Similar methods were employed to express AtPGP-1 in *Arabidopsis thaliana* plants with the following modifications. The AtPGP-1 coding region was subcloned into a pBIN vector lacking the GUS gene as described in Sidler, et al., 1998, *The Plant Cell* 10:1623-1636. This plasmid was then transformed into *A. tumefaciens* as described above, which was used to infect root calli to produce transgenic plants expressing AtPGP-1.

Plant growth: *Arabidopsis thaliana* seeds were sown in a solid germination media containing MS salt, 2% sucrose, 0.8% agar, and vitamins (Valvekens, D. et al., 1992, *Proc. Natl. Acad. Sci. USA* 85:5536-5540. For selection assays, cycloheximide was spread on the media to achieve a final concentration of 250 ng/ml. Plant growth was measured by germination percentage after 6-30 days.

RESULTS

Effect of over-expression of AtPGP-1 in yeast: When a yeast mutant, YMR4, which is deficient in two major extracellular phosphatases and tends to accumulate ATP extracellularly, was grown in a potent cellular toxin, cycloheximide, it did not grow whereas a wild-type yeast strain, INVSC1, did grow in the presence of cycloheximide (Figure 3A). Surprisingly, expression of the plant multidrug resistance (MDR) gene, AtPGP-1, enabled the yeast mutant to grow in the toxin (Figure 3B and Figure 5A below). The presence of AtPGP-1 in the wild-type yeast did not have any effect when grown in the presence of cycloheximide (Figure 3B). The same result was obtained when the yeast strains were cultured in nigericin (Figure 3C, 3D, Table 2, Figure 5B). In Figure 3C and 3D, starting from the top of the dish clockwise,

the cells are as follows: INVSC1 (wild-type) overexpressing AtPGP-1, YMR4 containing the vector alone, YMR4 overexpressing AtPGP-1, and INVSC1 containing the vector alone. When grown without drug, all the cells grow (Figure 3C). However, when grown in drug, only the YMR4 containing vector alone shows reduced growth. The survival of the AtPGP-1 transformed strains was due to the ability of the MDR1 channel to efflux the toxin, hence lowering the actual cellular concentration of the poison cycloheximide. The sensitivity of the untransformed mutant to the drug is likely due to a loss of the ATP gradient below a point at which endogenous transporters, similar to AtPGP-1 can function.

TABLE 2*

		Growth on Nigericin Drug selected cells			
		Day 0	Day 1	Day 2	Day 3
	ymr4mdr1	0	0.016	0.238	1.135
	ymr4 pvt	0	0.002	0.001	0.576
	invscmdr1	0	0.018	0.445	1.23
	invsc pvt	0	0.018	0.241	1.047

* Graph showing the growth turbidity of YMR4 yeast over-expressing the *Arabidopsis* plant ABC transporter AtPGP-1 grown in nigericin.

Effect of over-expression of AtPGP-1 in plants: The over-expression of AtPGP-1 was able to confer resistance to cycloheximide in plants (Figure 4A and 6) and to the cytokinin, N₆-(2-isopentenyl) adenine (2IP) (Figure 4B). These results had not been observed previously and in fact, the prior art actually teaches away from this finding suggesting that over-expression of plant AtPGP-1 is not involved in drug resistance. See Sidler, M. et al., 1998, *The Plant Cell* 10:1623-1636. Therefore, this result was particularly unexpected in plants. Additionally, since *Arabidopsis* plants overexpressing AtPGP-1 are able to grow in both cycloheximide and cytokinin, this suggests that the conference of drug resistance by AtPGP-1 is likely to be seen with other chemicals as well and is not an isolated phenomenon.

Effect of over-expression of apyrase on drug resistance in plants: Another unexpected result was obtained when the plant apyrase gene was over-expressed in plants. Over-expression of apyrase in plants resulted in the conference of resistance to cycloheximide (Figure 4A and 6).

(2-isopentenyl) adenine (Figure 4B). In fact, over-expression of apyrase is surprisingly able to raise the germination rate above the level obtained by the over-expression of the MDR gene AtPGP-1 (Figure 4A, 4B and 6). Just as under-expression of phosphatase activity in a yeast mutant lacking two potent extracellular phosphatases diminished its resistance to cycloheximide (Figure 3A), over-expression of a powerful extracellular ATP phosphatase in plants bolstered resistance. The fact that higher resistance was found in plants genetically manipulated only with respect to phosphatase over-expression and not MDR1, indicates that there likely exists other ATP-symporters used in detoxification in addition to MDR1. Minimally, the stronger ATP gradient set up by apyrase in the transgenic plants affects the kinetics of the wild-type MDR1.

EXAMPLE 3: ATP EFFLUX IN YEAST AND PLANTS OVEREXPRESSING AtPGP-1

MATERIALS AND METHODS

ATP collection: Yeast cells used in the luciferase assays were grown for two days and then transferred to fresh media at the time of the assay. From this time forward, the cells were kept at room temperature on a rotator. Every hour a 1 ml aliquot was taken, the cells in the aliquot were counted on a hemocytometer, a methylene blue viability assay was performed (Boyum, R. and Guidotti, G., 1997, *Microbiology* 143:1901-1908), the cells were centrifuged, and the supernatant was stored in liquid nitrogen until all the aliquots were collected. For luciferase assays involving plants, *Arabidopsis thaliana* plants were grown in sterile culture at 22° C under 150-200 μ E of continuous light for at least 15 days. Foliar ATP was collected by placing a single 30 μ l drop of luciferase buffer (Analytical Luminescence Laboratory, Cockeysville, Md.) on a leaf and, without making direct physical contact with the plant, the droplet was immediately collected and snap frozen. For each leaf, the area was approximated as an integrated area of a 2-D image of the leaf using NIH1.52 software (Shareware, NIH).

Luminometry: Samples were reconstituted to a 100 μ l final volume in Firelight™ buffer (Analytical Luminescence Laboratory, Cockeysville, MD). After the buffer was added, all samples were kept on ice. ATP standards were reconstituted in 100 μ l of Firelight™ buffer and the standards and sample were loaded into a 96-well plate and read on an automated Dynex Technologies Model MLX luminometer (Dynex Technologies, Chantilly, Va.). Samples were processed with the addition of 50 μ l of Firelight™ enzyme (Analytical Luminescence Laboratory, Cockeysville, MD) followed by a reading delay of 1.0 second and

an integration time of 10 seconds. Output was taken as an average for the integration time and then averaged for multiple samples. The sample handling time was less than 2 hours.

Pulse Chase experiments: Yeast were grown to saturation in liquid medium, as described above, centrifuged, and resuspended in fresh medium containing 1 $\mu\text{Ci/ml}$ ^3H -adenosine (Amersham, Arlington Heights, IL.). The cells were rotated at room temperature for 20 minutes to allow adenosine uptake. After 20 minutes the cells were centrifuged. The pellet was washed twice in ice cold medium, resuspended in culture medium at room temperature, divided equally between five types (five per cell line), and placed on a rotator. Every ten minutes a separate tube from each cell line was centrifuged and the pellet and supernatant were placed in separate scintillation vials. The efflux activity was expressed as the ratio of counts in the supernatant to counts in the pellet.

RESULTS

The ATP effluxed by the plant MDR1, AtPGP-1, over-expressed in yeast: In wild-type cells there is a steady-state level of ATP in the extracellular fluid, which is to say that the ATP outside the cells is rapidly degraded by phosphatases and does not accumulate over time (Figure 7). However, the expression of the AtPGP-1 doubled this steady-state level (Figure 8). If the yeast mutant, YMR4, which is deficient in extracellular phosphatase activity, is analyzed, there was a noticeable accumulation of ATP in the extracellular fluid compared to a control mutant transformed with empty plasmid pVT101 (Figure 9). In addition to ATP measurements based on luminometry performed on a kinetic time-scale of hours, an earlier differential ATP efflux in MDR1 expressing cells by pulse chase experiments was demonstrated (Figure 10). Furthermore, *Arabidopsis thaliana* plants from two independently transformed lines, that constitutively express the AtPGP-1 protein, showed a significant accumulation of ATP on their leaf surfaces (Figure 11). Taken together, these data demonstrate the absolute ability of plant MDR1, AtPGP-1, to transport ATP from inside the cell to the outside. Moreover, these data show that ATP efflux channels and phosphatases both have roles in the steady-state level of ATP outside of the cell. This is the first demonstration of the importance of extracellular ATP steady-state levels, and the importance of an ATP gradient across biological membranes in the modulation of drug resistance.

EXAMPLE 4: A TWO-COMPONENT SYSTEM IS FOUND IN ARABIDOPSIS PLANTS
MATERIALS AND METHODS

Plant Growth: *Arabidopsis* seeds were sown in a solid germination media containing MS salts (Sigma Chemical, St. Louis, Mo.), 2% sucrose, 0.8% agar, and vitamins (Valvekens, D. et al., 1992, *Proc. Natl. Acad. Sci. USA* 85:5536-5540). For selection assays, one of the following, or a combination of both, was added to media (cooled to less than 50°C before adding) immediately prior to pouring into plates: cycloheximide at a final concentration of 500 ng/ml; α,β -methyleneadenosine 5'-diphosphate at a final concentration of 1mM. Plant growth was measured by germination percentage after 10-20 days.

All other materials and methods were discussed above in Example 2.

RESULTS

Effects of phosphatase inhibitor on plants overexpressing AtPGP-1: Figure 12 shows that when wild-type and AtPGP-1 overexpressing (MDR OE) *Arabidopsis thaliana* plants were either treated with nothing (lane 1), cycloheximide (lane 2), α,β -methyleneadenosine 5'-diphosphate (phosphatase inhibitor) (lane 3), or cycloheximide and phosphatase inhibitor (lane 4), both the wild-type and the AtPGP-1 overexpressing plants were affected similarly by the presence of phosphatase inhibitor. While the AtPGP-1 overexpressing plants grew significantly better in the presence of cycloheximide alone with a 50% germination rate for the AtPGP-1 overexpressing plants and a 2% germination rate for the wild-type plants, similar germination rates were seen for both the AtPGP-1 overexpressing and wild-type plants in the presence of either phosphatase inhibitor alone (83% and 90% germination respectively) or cycloheximide plus phosphatase inhibitor (no germination at all). The addition of phosphatase inhibitor surprisingly destroys the ability of the AtPGP-expressing plants to grow in the presence of cycloheximide. These data suggest that phosphatases are involved in the conference of drug resistance in plants and that there is a two-component system similar to that demonstrated in yeast in Example 2 and 3 above in which an MDR-like protein and an ATP-gradient-maintaining ecto-phosphatase are important in modulating drug resistance.

EXAMPLE 5: THE ATP GRADIENT DIRECTLY EFFECTS DRUG RESISTANCE IN
CELLS

MATERIAL AND METHODS

Cell lines: Cell lines were the same as those described above in Example 2 and 3. YMR4

5 MDR1 is the phosphatase mutant yeast strain overexpressing AtPGP-1; YMR4 pVT101 contains vector alone; INVSC MDR1 is the wild-type yeast strain overexpressing AtPGP-1; and INVSC pVT101 contains vector alone.

Selection in drug: To create drug resistant yeast strains, all four cell lines were grown up in the presence of 500 ng/ml of cycloheximide, and transferred to other cycloheximide containing
10 plates after a period of four to six days. This transfer of cell lines and subculturing continued such that the yeast cells grew in the presence of cycloheximide for a period of at least a month.

Cells cultured in media alone: To create cell lines that had not been preselected for their ability to grow in drug, yeast strains were grown on plates containing YNB (Bio101, Vista, CA) without uracil (-URA) to maintain the presence of the vector (which supplies URA) without
15 any drugs added.

Growth of cells in suspension for ATP and drug selection experiments: Cells were transferred into 5 ml YNB -URA liquid media for turbidity measurements. All cell lines (both non-drug selected and drug-selected) were grown in media with the addition of either nothing, 500
20 ng/ml cycloheximide, 100 mM ATP, or 500 ng/ml cycloheximide and 100 mM ATP. Turbidity readings were taken after 48 hours.

Growth of cell lines in suspension for salvage pathway experiments: All cell lines were grown in liquid media either containing drug (for the drug selected lines) or not containing drug (for
25 the non-drug selected lines). When the cultures reached a turbidity of 1.00 as measured at a wavelength of 600 in a spectrophotometer ($OD_{600} = 1.00$), 10 μ l of each culture was then removed and placed in either media with nothing added, 3 mM potassium phosphate; 3 mM adenosine; 9 mM potassium phosphate and 3 mM adenosine (for controls); potassium phosphate and cycloheximide; adenosine and cycloheximide; adenosine, cycloheximide, and
30 potassium phosphate. Cell cultures were further grown for 72 hours, and their turbidity was determined by OD_{600} readings on a spectrophotometer.

Growth of cell lines for nigericin experiments: Drug selected lines were removed from cycloheximide containing plates and placed in 5 ml liquid media containing 5 ng/ml cycloheximide. Cell cultures were allowed to grow until they reached an OD₆₀₀ reading of 1.00, and then 10 µl from each culture was removed and transferred to culture tubes containing 5 ml of liquid media and 25 µg/ml nigericin. OD₆₀₀ readings were recorded daily for a period of up to 72 hours to determine growth.

RESULTS

An ATP gradient is critical in MDR: The importance of the ATP gradient in MDR in yeast cells was demonstrated by showing that the growth of cells which were previously grown in drug and had developed resistance to the drug, were not able to grow in high levels of ATP unless they were overexpressing AtPGP-1 (TABLE 3). Cells which had not been previously selected in drug were able to grow in the presence of high levels of ATP (TABLE 3). These data emphasize that the loss of an ATP gradient in previously resistant cell lines abolishes resistance. This result is new to the understanding of MDR and has led to vast insight into the understanding of the mechanism by which MDR-ABC transporters confer resistance to cells and to methods to modulate such resistance. Moreover, when cells were grown in high levels of ATP and drug (cycloheximide), even the cell lines which had previously showed resistance to drug were unable to grow in the presence of drug and ATP. These data indicate that when the ATP gradient across biological membranes is destroyed (by the presence of high extracellular levels of ATP), efflux of drugs cannot be achieved and therefore, drug resistance is abolished. In summary, the multi-drug resistance channel is not functional without an ATP gradient.

Table 3*

	Drug selected Cells	Cells cultured only on Media
<u>Cycloheximide</u>		
ymr4mdr1	0.745	0.014
ymr4 pvt	0.017	0.016
invscmdr1	0.683	0.013
invsc pvt	0.985	0.005

	Drug selected Cells	Cells cultured only on Media
<u>ATP+cycloheximide</u>		
ymr4mdr1	0.001	0.001
ymr4 pvt	0.002	0.001
invscmdr1	0.001	0.002
invsc pvt	0.001	0.002
<u>ATP</u>		
ymr4mdr1	0.016	0.585
ymr4 pvt	0.001	0.697
invscmdr1	0.271	1.267
invsc pvt	0.052	0.213
<u>Media alone</u>		
ymr4mdr1	1.477	1.478
ymr4 pvt	1.437	1.484
invscmdr1	1.498	1.483
invsc pvt	1.488	1.435

* Growth effects of cycloheximide and extracellular ATP on wild-type and MDR1 overexpressing *S. cerevisiae* yeast cells which have either never seen cycloheximide or which have been previously selected in cycloheximide.

The drug resistance is not due to an adenosine salvage pathway: In order to address whether the involvement of a nucleotide salvage pathway was responsible for the results of the present invention, yeast cells were cultured in the presence of extracellular adenosine and extracellular phosphate. The acid phosphatase yeast mutant, YMR4, was selected because its decreased ecto-phosphatase activity makes it an ideal candidate for studying the effect of extracellular nucleotides on growth. If an adenosine salvage pathway were involved, then the presence of extracellular adenosine or possibly phosphate should help cells recoup the intracellular ATP losses due to ATP/drug efflux and should help cells grow in the presence of drug whether or not the cells were overexpressing AtPGP-1. In contrast, however, the addition of adenosine or phosphate to the media did not enhance resistance to the cells (TABLE 4). In fact, cells overexpressing AtPGP-1 grew best in drug alone, with the addition of adenosine and/or phosphate being slightly inhibitory. Furthermore, cells which did not express AtPGP-1 were unable to grow in drug regardless of the presence of adenosine and/or phosphate. These data

suggest that an adenosine salvage pathway is not the principal mechanism at work in the present invention.

Table 4*

5	<u>Media alone</u>	<u>Cycloheximide</u>
	ymr4 mdr1 1.376	ymr4 mdr1 0.937
	ymr4 pvt 1.429	ymr4 pvt 0.001
10	<u>PO₄ alone</u>	<u>PO₄ and Cycloheximide</u>
	ymr4 mdr1 1.351	ymr4 mdr1 0.541
	ymr4 pvt 1.341	ymr4 pvt 0.001
15	<u>Adenosine alone</u>	<u>Adenosine and Cycloheximide</u>
	ymr4 mdr1 1.319	ymr4 mdr1 0.632
	ymr4 pvt 1.354	ymr4 pvt 0.002
	<u>Adenosine and PO₄ alone</u>	<u>Adenosine and PO₄ and Cycloheximide</u>
	ymr4 mdr1 0.899	ymr4 mdr1 0.389
	ymr4 pvt 1.342	ymr4 pvt 0.001

* Growth effects of cycloheximide, adenosine and phosphate on wild-type and AtPGP-1 overexpressing *S. cerevisiae* yeast cells.

20 EXAMPLE 6: HIGH THROUGHPUT SCREEN FOR ISOLATING APYRASE INHIBITORS

MATERIAL AND METHODS

Small Molecule Library: A small molecule library (DIVERSet format F), which was specifically constructed to maximize structural diversity in a relatively small library (9600
25 compounds), was obtained from ChemBridge Corporation (San Diego, CA). The small molecules (supplied in 0.1 mg dehydrated aliquots) were dissolved in DMSO, transferred to a 96 well plate, and tested for their ability to inhibit apyrase activity.

The assay: A stringent screen to test the ability of small molecules to disrupt the ATPase activity of the apyrase enzyme was developed based on phosphate-mobylate complexation.
30 The assay was a modification of a phospholipase assay developed by Hergenrother et al.

(*Lipids* 32:783-788 (1997)). Under normal conditions, the apyrase enzyme liberates phosphate from ATP present in the reaction. The liberated phosphate quickly forms a complex upon addition of a small amount of acidified molybdate and ascorbate allowing for the production of a very dark blue color (the less phosphate liberated, the less blue color).

5 Control reactions were performed with heat inactivated apyrase enzyme. Color intensity was detected on an Alpha Imager 2000 with AlphaEase™ software (Alpha Innotech, San Leandro, CA). Color changes were also evident by the naked eye. A Biomek 2000 robot (Beckman, Fullerton, CA) was used for screening the 9600 samples.

10 To each well of the 96 well plates containing a small molecule from the library, 100 µl of reaction buffer (60 mM HEPES, 3 mM MgCl₂, 3 mM CaCl₂, 3 mM ATP pH 7.0) was added. The apyrase (potato apyrase grade VI, Sigma Chemical, St. Louis, MO) enzyme (0.1 units) was added in a 5 µl volume and the reaction was allowed to proceed at room temperature for 60 minutes.

Three buffers were used to visualize activity:

15 Buffer A: 2% Ammonium molybdate in water

Buffer B: 11% Ascorbic acid in 37.5% aqueous TCA.

Buffer C: 2% trisodium citrate, 2% acetic acid.

Immediately before developing the assay, buffers A and B were mixed in a 1:1.5 ratio. 50 µl of A:B was added to each well. The 96 well plate was then vibrated on a
20 table surface to mix the solution. The deep blue color developed after approximately 2 minutes. After 2 minutes, 50 µl of buffer C was added to each well and the blue color became darker, increasing the sensitivity of the assay. The color intensified for up to one hour with no accompanying color change in the control wells containing heat inactivated apyrase enzyme. The color intensity for a single plate was measured on an Alpha Imager 2000 with
25 AlphaEase™ software (Alpha Innotech, San Leandro, CA).

RESULTS:

Nineteen positives were identified from the 9600 compound DIVERSet library. Dose response assays revealed that fourteen showed weak inhibition, two showed medium inhibition (Formulas IV and V), and three showed relatively strong inhibition (Formulas I, II and III).

30

WHAT IS CLAIMED IS:

1. A method for increasing or decreasing drug resistance in a target bacteria, yeast, plant or mammalian cell comprising altering the ATP gradient across the biological membrane of the target cell.
- 5 2. A method for altering the ATP gradient across the biological membrane of a target bacteria, yeast, plant or mammalian cell to achieve an increase in drug resistance comprising up-regulating an ecto-phosphatase in the target cell.
3. The method of claim 2 further comprising up-regulating an ABC transporter in the target cell.
- 10 4. A method for altering the ATP gradient across the biological membrane of a target bacteria, yeast, plant or mammalian cell to achieve a decrease in drug resistance comprising down-regulating an ecto-phosphatase in a target cell.
5. The method of claim 4 further comprising down-regulating an ABC transporter in a target cell.
- 15 6. A method for altering the ATP gradient across the biological membrane of a plant cell to achieve an increase in drug resistance comprising up-regulating an ABC transporter in the target cell.
7. A method for altering the ATP gradient across the biological membrane of a plant cell to achieve a decrease in drug resistance comprising down-regulating an ABC
- 20 transporter in the target cell.
8. A method for augmenting the chemotherapeutic effectiveness of a chemotherapeutic molecule by decreasing resistance to the chemotherapeutic molecule in a target cell comprising down-regulating an ecto-phosphatase in a target cell.
9. The method of claim 8 further comprising down-regulating an ABC transporter in
- 25 the target cell.
10. A method for conferring herbicide resistance to a plant comprising up-regulating an ecto-phosphatase in the target cell.
11. The method of claim 10 further comprising up-regulating an ABC transporter in
- 30 the target cell.


12. A method for increasing sensitivity to a drug molecule to inhibit or ameliorate microorganism infection in animals and humans by altering the ATP gradient across the biological membrane of a microorganism to achieve a decrease in drug resistance comprising down-regulating an ecto-phosphatase in a target cell.

5 13. The method of claim 12 further comprising down-regulating an ABC transporter in a cell of the microorganism.


14. The method of claim 2, 4, 8, 10 or 12 wherein the ecto-phosphatase is selected from the group consisting of *Pisum sativum* apyrase (GenBank accession # Z32743) and *Homo sapiens* apyrases (GenBank accession # AF034840, AF039916, AF039917, 10 AF039918 and HSU87967).

15. The method of claim 3, 5, 6, 7, 9, 11 or 13 wherein the ABC transporter is selected from the group consisting of *Arabidopsis thaliana* AtPGP-1 (GenBank accession # X61370), *Homo sapiens* Pgp (GenBank accession # M29432), *Homo sapiens* MDR- β (PCT publication WO 98/46736), *Saccharomyces cerevisiae* STS1 (GenBank accession # X75916), 15 *Saccharomyces cerevisiae* Pdr5p (GenBank accession # 1420383), *Aspergillus fumigatus* Afu-MDR1 (U.S. Patent No. 5,705,352) and *Lactococcus lactis* LmrA (GenBank accession # U63741).

16. A method for inhibiting an ecto-phosphatase comprising contacting an ecto-phosphatase with an ecto-phosphatase inhibiting amount of an ecto-phosphatase inhibitor 20 selected from the group consisting of molecules having the Formulae I through XIX:



I




II


56

5

III



10




IV

15


20

V



25


VI



30


57

5




VII

10



VIII

15




IX

20

25

X


30



58


XI

5



XII


10



15


XIII

20




XIV

25




XV

30




59


XVI




XVII



XVIII




XIX




17. The method of claim 16 wherein the ecto-phosphatase inhibitory molecule is selected from the group consisting of molecules having the Formulae I through V:

5




10



15


III


20



25


30






5

18. The method of claim 16 wherein the ecto-phosphatase inhibitory molecule is selected from the group consisting of molecules having the Formulae I through III:



10


II



15

20

III



25


19. The method of claim 16 wherein the ecto-phosphatase is selected from the group consisting of *Pisum sativum* apyrase (GenBank accession # Z32743) and *Homo sapiens* apyrases (GenBank accession # AF034840, AF0399ecto-phosphatase, AF039917, AF039918 and HSU87967).

30


20. A method for decreasing drug resistance in a target bacteria, yeast, plant or mammalian cell comprising introducing to cells a drug resistance-inhibiting amount of an ecto-phosphatase inhibitory molecule selected from the group consisting of molecules having the Formulae I through XIX:

5

I



10




15

II


20

III







25


30




IV



5




10



15


20



25


XII

30




65

XIII




5



XIV


10



XV

15


20



XVI


25

30




XVII

XVIII



XIX



21. The method according to claim 20 wherein the mammalian cells are tumor cells.

22. A method of identifying an inhibitor of an ecto-phosphatase comprising

a) contacting the ecto-phosphatase with a small molecule in the presence of ATP under conditions wherein the ecto-phosphatase has ATPase activity,




b) incubating the ecto-phosphatase, small molecule and ATP for a period of time sufficient to liberate phosphate from the ATP, and

5 c) adding ammonium molybdate and ascorbic acid to the ecto-phosphatase, small molecule and ATP to form a complex with liberated phosphate and to generate a dark blue color, wherein inhibition of the ecto-phosphatase by the small molecule results in less phosphate liberated and less blue color.

10 23. The method according to claim 22 further comprising adding trisodium citrate and acetic acid.

24. The method according to claim 22 wherein the ecto-phosphatase is selected from the group consisting of *Pisum sativum* apyrase (GenBank accession # Z32743), *Homo sapiens* apyrase (GenBank accession # AF034840, AF0399ecto-phosphatase, AF039917, AF039918 and HSU87967) and potato apyrase (GenBank accession U58597).

15



SUBSTITUTE SHEET (RULE 26)

2/11


 ^3H ATP Uptake

FIG. 2A


 α ^{32}P ATP Uptake

FIG. 2B




 γ ^{32}P ATP Uptake

FIG. 2C

SUBSTITUTE SHEET (RULE 26)

3/11



4/11




FIG. 4A

5/11




FIG. 4B-2




FIG. 4B-3




FIG. 4B-1

6/11




FIG. 5A




FIG. 5B

7/11




FIG. 6




FIG. 7

SUBSTITUTE SHEET (RULE 26)

8/11




FIG. 8




FIG. 9

9/11




FIG. 10

Extracellular ^3H -adenosine (c.p.m.)

Intracellular ^3H -adenosine (c.p.m.)

10/11




FIG. 11

11/11





FIG. 12

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/05315

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/4, 7.21, 7.23, 7.32, 7.93, 468, 471, 483; 800/8, 14, 21, 278, 295, 298, 300 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) West, Agricola, Caplus, Biosis														
C. DOCUMENTS CONSIDERED TO BE RELEVANT														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
X --- Y	DECOTTIGNIES et al. ATPase and Multidrug Transport Activities of the Overexpressed Yeast ABC Protein Yor1p. J. Biol. Chem. 15 May 1998, Vol. 273, No. 20, pages 12612-12622, especially pages 12617-12619.	1-3, 6 ----- 5, 7, 15												
Y	DUDLER et al. Structure of an <i>mdr</i> -like Gene from <i>Arabidopsis thaliana</i> . J. Biol. Chem. 25 March 1992, Vol. 267, No. 9, pages 5882-5888, especially pages 5883-5885.	1-7, 11, 15												
X --- Y	GRANT et al. Overexpression Of Multidrug Resistance-associated Protein (MRP) Increases Resistance To Natural Product Drugs. Cancer Research. 15 January 1994, Vol. 54, pages 357-361, especially pages 358-360.	1 ----- 3-8, 15												
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"> <tr> <td>* Special categories of cited documents:</td> <td>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>*A* document defining the general state of the art which is not considered to be of particular relevance</td> <td>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>*E* earlier document published on or after the international filing date</td> <td>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>*Z* document member of the same patent family</td> </tr> <tr> <td>*O* document referring to an oral disclosure, use, exhibition or other means</td> <td></td> </tr> <tr> <td>*P* document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means		*P* document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family													
O document referring to an oral disclosure, use, exhibition or other means														
P document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 05 MAY 2000		Date of mailing of the international search report 26 May 2000 (26.05.00)												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer ASHWIN MEHTA  Telephone No. (703) 308-0196												

Form PCT/ISA/210 (second sheet) (July 1998)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/05315

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LU et al. AtMRP2, An Arabidopsis ATP Binding Cassette Transporter Able To Transport Glutathione S-Conjugates And Chlorophyll Catabolites: Functional Comparisons With AtMRP1. The Plant Cell. February 1998, Vol. 10, pages 267-282, especially pages 271-274.	1, 3-7, 10, 11, 14, 15
Y	KIBA et al. Specific Inhibition Of Cell Wall-Bound ATPases By Fungal Suppressor From <i>Mycosphaerella pinodes</i> . Plant Cell Physiol. 1995, Vol. 36, No. 5, pages 809-817, especially pages 813-814.	16-20, 22-24
Y	THOMAS et al. Apyrase Functions In Plant Phosphate Nutrition And Mobilizes Phosphate From Extracellular ATP. Plant Physiol. February 1999, Vol. 119, pages 543-551, see whole document.	1-5, 10, 11, 14, 15, 16-24
Y	WANG et al. CD39 Is An Ecto-(Ca ²⁺ , Mg ²⁺)-Apyrase. J. Biol. Chem. 26 April 1996, Vol. 271, No. 17, pages 9898-9901, especially pages 9899-9901.	1-5, 14, 15
Y	SIDLER et al. Involvement Of An ABC Transporter In A Developmental Pathway Regulating Hypocotyl Cell Elongation In The Light. The Plant Cell. October 1998, Vol. 10, No. 10, pages 1632-1636, see whole document.	1, 3, 5-7, 15

Form PCT/ISA/210 (continuation of second sheet) (July 1998)*

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/05315

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-7, 10, 11, 14, 15, 16-24
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1)) (July 1998)★

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/05315

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

C12N 5/04, 5/06, 1/16, 1/20, 15/67, 15/81, 15/82, 15/90; A01H 1/00, 5/00

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/4, 7.21, 7.23, 7.32, 7.93, 468, 471, 483; 800/8, 14, 21, 278, 295, 298, 300

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-7, 10, 11, 14 and 15, drawn to a first method for increasing or decreasing drug resistance in a target bacteria, plant, yeast or mammalian cell comprising altering the ATP gradient across the biological membrane of the cell.

Group II, claims 8 and 9, drawn to a second method, for augmenting the chemotherapeutic effectiveness of a chemotherapeutic molecule.

Group III, claims 12 and 13, drawn to a third method, for increasing sensitivity to a drug molecule to inhibit microorganism infection in animals and humans.

Group IV, claims 16-24, drawn to a fourth method, for inhibiting an ecto-phosphatase comprising contacting an ectophosphatase with an ecto-phosphatase inhibitor, and for identifying an ecto-phosphatase inhibitor.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The method for increasing drug resistance of Group I is not shared by the method for augmenting chemotherapeutic effectiveness of Group II, or with the method for inhibiting an ecto-phosphatase or for identifying an ecto-phosphatase inhibitor of Group IV. The plants of Group I are not shared with the methods of Groups II-IV. The chemotherapeutic molecules of Group II are not shared with the other groups. The microorganism infection of Group III is not shared with the other groups. The identification of ecto-phosphatase inhibitors of Group IV is not shared with the other groups.